

09/911, 072

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup> :</b> <b>C07D 239/42, 239/47, 239/52</b> <b>C07C 309/89, A01N 43/54</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 91/15478</b> <b>(43) International Publication Date:</b> <b>17 October 1991 (17.10.91)</b>
<b>(21) International Application Number:</b> <b>PCT/US91/01075</b> <b>(22) International Filing Date:</b> <b>26 February 1991 (26.02.91)</b>  <b>(30) Priority data:</b> <b>503,182</b> <b>2 April 1990 (02.04.90)</b> <b>US</b>  <b>(60) Parent Application or Grant</b> <b>(63) Related by Continuation</b> <b>US</b> <b>503,182 (CIP)</b> <b>Filed on</b> <b>2 April 1990 (02.04.90)</b>  <b>(71) Applicant (for all designated States except US):</b> <b>E.I. DU PONT DE NEMOURS AND COMPANY [US/US];</b> <b>1007 Market Street, Wilmington, DE 19898 (US).</b>		<b>(72) Inventor; and</b> <b>(75) Inventor/Applicant (for US only) :</b> <b>MOON, Marcus, P. [US/US];</b> <b>1703 North Union Street, Wilmington, DE 19806 (US).</b>  <b>(74) Agents:</b> <b>FISHER, Lynn, N. et al.; E.I. du Pont de Nemours and Company, Legal/Patent Records Center,</b> <b>1007 Market Street, Wilmington, DE 19898 (US).</b>  <b>(81) Designated States:</b> <b>AT (European patent), BE (European patent), BR, CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent), US.</b>  <b>Published</b> <b>With international search report.</b>
<b>(54) Title: CROP-SELECTIVE HERBICIDAL SULFONAMIDES</b>  <b>(57) Abstract</b> <p>This invention relates to novel sulfonamides and their use as crop selective herbicides. Compounds of the instant invention have demonstrated excellent control of weeds coupled with corn safety.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark				

**TITLE****CROP-SELECTIVE HERBICIDAL SULFONAMIDES**

5

**CROSS-REFERENCE TO RELATED APPLICATION**

This is a continuation-in-part of application  
Serial No. 07/503,182, filed April 4, 1990.


**BACKGROUND OF THE INVENTION**

This invention relates to novel sulfonamides  
10 and their use as crop selective herbicides.  
Compounds of the instant invention have demonstrated  
excellent control of weeds coupled with corn safety.

EP-A-120,814 discloses herbicidal  
sulfonylureas of the formula

15

20



25 wherein:

A is C<sub>1</sub>-C<sub>6</sub> haloalkyl;

R<sup>1</sup> is H, halogen, NO<sub>2</sub>, CN, C<sub>1</sub>-C<sub>4</sub> alkyl,  
C<sub>1</sub>-C<sub>4</sub> haloalkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub>  
alkylthio, C<sub>1</sub>-C<sub>4</sub> alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>  
30 alkylsulfonyl, COR<sup>6</sup>, NR<sup>7</sup>R<sup>8</sup>, CONR<sup>9</sup>R<sup>10</sup> or  
SO<sub>2</sub>NR<sup>11</sup>R<sup>12</sup>; and

R<sup>6</sup> is C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkoxy,  
C<sub>1</sub>-C<sub>4</sub> alkylthio, C<sub>2</sub>-C<sub>6</sub> alkoxyalkoxy,  
hydrogen, C<sub>1</sub>-C<sub>4</sub> alkyl or C<sub>1</sub>-C<sub>4</sub> haloalkyl.

35 Although this reference broadly embraces  
compounds of the instant invention, it does not teach  
or suggest these particular compounds or their  
herbicidal utility.


Additionally, a need still exists for herbicides because of world-wide food shortages. In addition, herbicides which are selective to important crops such as corn are particularly necessary. According to the instant invention, such compounds have been found.

# SUMMARY OF THE INVENTION

More specifically, this invention comprises novel compounds of Formula I, agriculturally suitable compositions containing them, and their method-of-use as preemergence and/or postemergence herbicides or plant growth regulants:

15

20



I

25

wherein:

- A is  $OR^1$  or  $N(CH_3)_2$ ;
  - $R^1$  is  $C_1-C_3$  alkyl,  $CH_2CH=CH_2$ ,  $CH_2C\equiv CH$ ,  $CH_2CH_2Cl$  or  $CH_2CH_2OCH_3$ ;
  - $R^2$  is  $CH_2F$ ,  $CHF_2$ ,  $CHFCH_3$  or  $CH_2CN$ ;
  - X is  $CH_3$  or  $OCH_3$ ; and
  - Y is H, Cl,  $CH_3$ ,  $C_2H_5$ ,  $OCH_3$  or  $OCF_2H$ ;
- and their agriculturally suitable salts; provided that when Y is Cl, then X is  $OCH_3$ .

35

In the above definitions, the term "alkyl" denotes straight chain or branched alkyl, e.g.

5 methyl, ethyl, n-propyl or isopropyl. The total number of carbon atoms in a substituent group is indicated by the C<sub>i</sub>-C<sub>j</sub> prefix where i and j are numbers 1 to 3. For example, C<sub>1</sub>-C<sub>3</sub> alkyl would designate methyl through propyl (both n-propyl and isopropyl).

10 Preferred for reasons of increased ease of synthesis and/or greater herbicidal efficacy and/or crop safety are:

1. Compounds of Formula I wherein  
A is OR<sup>1</sup>; and  
R<sup>1</sup> is CH<sub>3</sub>, CH<sub>2</sub>CH<sub>3</sub> or CH(CH<sub>3</sub>)<sub>2</sub>.
- 15 2. Compounds of Preferred 1 where  
when one of X and Y is CH<sub>3</sub>, then the other of X and Y is other than OCH<sub>3</sub>.

Specifically preferred for reason of greatest safety to corn (maize) is:

- 20 • Methyl 3-(cyanomethyl)-2-[[[(4,6-dimethyl-2-pyrimidinyl)amino]carbonyl]-amino]sulfonyl]benzoate (Formula I: A is OR<sup>1</sup>, R<sup>1</sup> is CH<sub>3</sub>, R<sup>2</sup> is CH<sub>2</sub>CN, X and Y are CH<sub>3</sub>).

25 Specifically Preferred for reason of greatest safety to corn, wheat and barley are:

- 30 • Methyl 2-[[[(4-chloro-6-methoxy-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]-3-(cyanomethyl)benzoate (Formula I: A is OR<sup>1</sup>, R<sup>1</sup> is CH<sub>3</sub>, R<sup>2</sup> is CH<sub>2</sub>CN, X is OCH<sub>3</sub>, Y is Cl);
- Methyl 3-(cyanomethyl)-2-[[[(4-(difluoromethoxy)-6-methoxy-2-pyrimidinyl)-amino]carbonyl]amino]sulfonyl]benzoate (Formula I: A is OR<sup>1</sup>, R<sup>1</sup> is CH<sub>3</sub>, R<sup>2</sup> is CH<sub>2</sub>CN, X is OCH<sub>3</sub>, Y is OCF<sub>2</sub>H);
- 35 • Methyl 3-(difluoromethyl)-2-[[[(4-methoxy-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]-

benzoate (Formula I: A is  $OR^1$ ,  $R^1$  is  $CH_3$ ,  $R^2$  is  $CHF_2$ , X is  $OCH_3$ , Y is H);

- 5 • Methyl 3-(fluoromethyl)-2-[[[(4-methoxy-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]-benzoate (Formula I: A is  $OR^1$ ,  $R^1$  is  $CH_3$ ,  $R^2$  is  $CH_2F$ , X is  $OCH_3$ , Y is H).

Specifically preferred for reason of greatest  
10 safety to wheat and barley is:


- Methyl 2-[[[(4,6-dimethoxy-2-pyrimidinyl)-amino]carbonyl]amino]sulfonyl]-3-(fluoromethyl)benzoate (Formula I: A is  $OR^1$ ,  $R^1$  is  $CH_3$ ,  $R^2$  is  $CH_2F$ , X and Y are  $OCH_3$ ).
- 15

Specifically preferred for reason of greatest  
safety to corn, wheat, barley and rice is:

- Ethyl 2-[[[(4-chloro-6-methoxy-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]-3-(fluoromethyl)benzoate (Formula I: A is  $OR^1$ ,  $R^1$  is  $CH_2CH_3$ ,  $R^2$  is  $CH_2F$ , X is  $OCH_3$ , Y is Cl).
- 20

This invention also comprises novel compounds,  
such as the sulfonyl benzoates of Formula II, useful  
25 as intermediates for preparation of the compounds of  
Formula I:

30



35

II

wherein:

- 5           A is  $\text{OR}^1$  or  $\text{N}(\text{CH}_3)_2$ ;  
            $\text{R}^1$  is  $\text{C}_1\text{-C}_3$  alkyl,  $\text{CH}_2\text{CH}=\text{CH}_2$ ,  $\text{CH}_2\text{C}\equiv\text{CH}$ ,  
            $\text{CH}_2\text{CH}_2\text{Cl}$  or  $\text{CH}_2\text{CH}_2\text{OCH}_3$ ;  
            $\text{R}^2$  is  $\text{CH}_2\text{F}$ ,  $\text{CHF}_2$ ,  $\text{CHFCH}_3$  or  $\text{CH}_2\text{CN}$ ; and  
            $\text{Z}^1$  is  $\text{Cl}$  or  $\text{NHSiR}^3\text{R}^4\text{R}^5$ ;  
            $\text{R}^3$  is  $\text{C}_1\text{-C}_4$  alkyl;  
 10           $\text{R}^4$  is  $\text{C}_1\text{-C}_4$  alkyl; and  
            $\text{R}^5$  is  $\text{C}_1\text{-C}_4$  alkyl.


Preferred for reason of increased herbicidal activity of final products of Formula I, are intermediates of Formula II wherein A is  $\text{OR}^1$  and  $\text{R}^1$  is  $\text{C}_1\text{-C}_2$  alkyl.

Preferred for reason of increased ease of synthesis are intermediates of the above preferred wherein  $\text{R}^3$  and  $\text{R}^4$  are  $\text{CH}_3$  and  $\text{R}^5$  is  $\text{C}(\text{CH}_3)_3$ .

#### DETAILED DESCRIPTION OF THE INVENTION

20

25



(1)

30

#### SYNTHESIS


The compounds of Formula (1), which corresponds to Formula I, can be prepared by one or more of the methods described below. The proper choice of reaction sequences for a given compound will be known to one skilled in the art.

As shown in Equation 1, many of the compounds of Formula (1) are prepared by reacting a silyl sulfonamide of Formula (2) with a pyrimidine carbamate of Formula (3).  $R^3$ ,  $R^4$ , and  $R^5$  are independently  $C_1$  to  $C_4$  alkyl.

Equation 1

10

15



20

The reaction is carried out at 0°C to 50°C in a solvent such as acetonitrile, dioxane, or tetrahydrofuran; in the presence of a fluoride ion source such as cesium fluoride, or tetrabutylammonium fluoride for 0.1 to 2 hours. A catalytic amount of base, such as 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU), increases the reaction rate.

25

Alternatively, some of the sulfonamides of Formula (4) can be prepared and reacted with pyrimidine carbamates of Formula (3) to give compounds of Formula (1) as shown in Equation 2. The reaction is carried out at 0°C to 50°C in a solvent, such as acetonitrile, dioxane, or tetrahydrofuran, in the presence of a non-nucleophilic base, such as DBU for 0.2 to 2 hours. U.S. Patent 4,604,131 discloses details for similar reactions and is herein incorporated by reference.


30

35



Equation 2

5




15

Sulfonyl chlorides of Formula (5) may be reacted with cyanate anion in the presence of pyrimidine amines of Formula (6) to give compounds of Formula (1) as shown in Equation 3.

20 Equation 3

25



30

The reaction is carried out by mixing one equivalent of sulfonyl chloride (5), pyrimidine amine (6), a metal cyanate, such as potassium cyanate, and a catalytic amount of an amin bas , such as Aliquat 336 (Tricaprylylmethylammonium chlorid ), in a solvent such as acetonitrile for 0.2 to 10 days.


35

Alternatively, one can use a tetraalkylammonium cyanate, such as tetraethylammonium cyanate, to effect this reaction by the method described in U.S. Patent 4,604,131 herein incorporated by reference.

Silyl sulfonamides of Formula (2) are prepared by reacting sulfonyl chlorides of Formula (5) with a trialkylsilyl amine of Formula (7), such as t-butyldimethylsilyl amine, as shown in Equation 4. Equation 4

15

20



The reaction is carried out at 0°C to 30°C in a solvent, such as dichloromethane, in the presence of 1 to 2 equivalents of the amine and 1 equivalent of a bicarbonate. J. R. Bowser et. al. describe methods to prepare silyl amines of Formula (7) in Inorganic Chemistry 17, 1882 (1978).


The sulfonyl chlorides of Formula (5) and the silyl sulfonamides of Formula (2) correspond to the intermediates of Formula II.

Sulfonyl chlorides of Formula (5) can be reacted with no more than 2 equivalents of ammonia or, alternatively, 1 equivalent of ammonia and 1 equivalent of a base to give sulfonamides of Formula (4) as outlined in Equation 5.

Equation 5

5

10



15 Sulfonamides of Formula (4) tend to be unstable and can be difficult to isolate.


Sulfonyl chlorides of Formula (5) are prepared by the methods shown in Equations 6 and 7.

20 Sulfur-containing compounds of Formula (8) are oxidized with chlorine as shown in Equation 6.  $\text{R}^6$  is H, alkyl, benzyl or carbamoyl.

Equation 6

25

30



35

The reaction of Equation 6 is carried out by contacting compounds of Formula (8) in a solvent, such as acetic acid or propionic acid, with at least


3.0 equivalents of chlorine in the presence of at least 2.5 equivalents of water for 0.2 to 5 hours at -20°C to 30°C. A. Wagenaar teaches specific reaction conditions for related compounds in *Recl. Trav. Chim. Pays-Bas* 101, 91 (1982).

Alternatively, reaction of compounds of Formula (8), where R<sup>6</sup> is H or benzyl, with a hypochlorite solution, such as 5% NaOCl, can provide sulfonyl chlorides of Formula (5). Reaction conditions for similar reactions are obvious to one skilled in the art and are described in EP-A-142,152.

As shown in Equation 7, below, sulfonyl chlorides of Formula (5) may be prepared from the corresponding anilines of Formula (9) by a Meerwein reaction.

Equation 7

20



30 The aniline is diazotized and then reacted with sulfur dioxide and cupric chloride analogous to the teachings of Yale and Sowinski, *J. Org. Chem.* 25, 1824 (1960). Alternatively, the hydrochloride salts of anilines of Formula (9) can be diazotized in an organic solvent with an alkyl nitrite and react d  
35 with sulfur dioxide to give sulfonyl chlorides (5) analogous to the teachings of M. Doyl, *J. Org. Chem.* 42, 2426, 2431 (1977).

The pyrimidine carbamates of Formula (3) and the pyrimidine amines of Formula (6) are prepared by the methods described and referenced in EP-A-72,347, EP-A-164,269, EP-A-173,498, U.S. Patent 4,540,782, and U.S. Patent 4,666,506, herein incorporated by reference.

Agriculturally suitable salts of compounds of Formula I are also useful herbicides and can be prepared in a number of ways known to the art. For example, metal salts can be made by contacting compounds of Formula I with a solution of an alkali or alkaline earth metal salt having a sufficiently basic anion (e.g., hydroxide, alkoxide, carbonate or hydroxide). Quaternary amine salts can be made by similar techniques.

Salts of compounds of Formula I can also be prepared by exchange of one cation for another. Cationic exchange can be effected by direct contact of an aqueous solution of a salt of a compound of Formula I (e.g., alkali or quaternary amine salt) with a solution containing the cation to be exchanged. This method is most effective when the desired salt containing the exchanged cation is insoluble in water and can be separated by filtration.

Exchange may also be effected by passing an aqueous solution of a salt of a compound of Formula I (e.g., an alkali metal or quaternary amine salt) through a column packed with a cation exchange resin containing the cation to be exchanged for that of the original salt and the desired product is eluted from the column. This method is particularly useful when the desired salt is water-soluble, e.g., a potassium, sodium or calcium salt.

Acid addition salts, useful in this invention, can be obtained by reacting a compound of Formula I

with a suitable acid, .g., p-toluenesulfonic acid, trichloroacetic acid or the like.

5           The preparation of the compounds of this invention is further illustrated by the following specific examples. Temperatures are reported in degrees Celsius; abbreviations for nuclear magnetic resonance (NMR) are: s = singlet, d = doublet, t =  
10   triplet, m = multiplet, and peak positions are reported as parts per million downfield from internal tetramethylsilane. Infrared (IR) peak positions are given in reciprocal centimeters ( $\text{cm}^{-1}$ ) and sh denotes a shoulder.

15

#### EXAMPLE 1

##### Methyl 3-(Cyanomethyl)-2-nitrobenzoate

To a stirred solution of potassium cyanide (6.2 g) in methanol (100 mL) and water (150 mL) at 0°C was added an acetonitrile solution of methyl  
20   3-(bromomethyl)-2-nitrobenzoate (24.9 g). A catalytic amount of 18-crown-6 ether was then added. After stirring overnight at room temperature, the reaction was extracted with ethyl acetate. The combined extracts were washed with a brine solution,  
25   dried over magnesium sulfate, filtered, concentrated, and chromatographed on silica gel eluting with 20% ethyl acetate in hexanes to give 8.5 g of the title compound as a white solid.

90 MHz NMR ( $\text{CDCl}_3$ )  $\delta$ :   3.83 (s, 2H,  $\text{CH}_2$ );  
30                           3.93 (s, 3H,  $\text{CH}_3$ ); and  
                              7.9 (m, 3H, arom.)

#### EXAMPLE 2

##### 2-(Methoxycarbonyl)-6-(cyanomethyl)-benzenesulfonyl Chloride

35           To a stirred suspension of dry tetrahydrofuran (150 mL) and 26% potassium hydride in oil (6.66 g) was slowly added benzyl mercaptan (4.82 mL) at 0°C under a nitrogen atmosphere. After 15 minutes, methyl

3-(cyanomethyl)-2-nitrobenzoate (8.5 g) was added and the resulting mixture was stirred at room temperature overnight. The reaction was contacted with 6N sodium hydroxide (25 mL) and extracted with ethyl acetate. The extracts were washed with a brine solution, dried over magnesium sulfate, filtered, and concentrated to give 9.0 g of crude methyl 3-(cyanomethyl)-2-(phenylmethylthio)benzoate as a purple oil.

The crude thioether was stirred in a mixture of dichloromethane (200 mL) and 6N hydrochloric acid (25 mL) at 0°C, as 5% sodium hypochlorite (180 mL) was slowly added. The reaction mixture was then stirred at 0°C for 2 hours. The dichloromethane layer was washed with water, dried over magnesium sulfate, filtered, concentrated, and chromatographed on silica gel eluting with 25% ethyl acetate in hexanes to give 1.7 g of the title compound as a yellow solid.

90 MHz NMR (CDCl<sub>3</sub>) δ: 3.98 (s, 3H, CH<sub>3</sub>);  
4.41 (s, 2H, CH<sub>2</sub>); and  
7.83 (m, 3H, arom.).

### EXAMPLE 3

Methyl 3-(Cyanomethyl)-2-[[[(1,1-dimethylethyl)-dimethylsilyl]aminol]sulfonyl]benzoate

A suspension of 2-(methoxycarbonyl)-6-(cyanomethyl)benzenesulfonyl chloride (10.7 g), amino-t-butyltrimethylsilane (10.3 g), and sodium bicarbonate (3.36 g) in dichloromethane (200 mL) was stirred at room temperature in a stoppered flask for 4 days. The organic phase was washed with water, dried over magnesium sulfate, filtered, concentrated, and chromatographed on silica gel eluting with 30% ethyl acetate in hexanes to give 5.0 g of the title compound as a yellow solid, m.p. 113-115°C.

90 MHz NMR (CDCl<sub>3</sub>) δ: 0.32 (s, 6H, SiCH<sub>3</sub>);  
0.95 (s, 9H, C(CH<sub>3</sub>)<sub>3</sub>);

14

5  
3.99 (s, 3H, OCH<sub>3</sub>);  
4.42 (s, 2H, CH<sub>2</sub>CN);  
5.91 (s, 1H, NH);  
7.6 (m, 2H, arom.); and  
7.8 (m, 1H, arom.).

IR (mineral oil) 3290, 2250, and 1710 cm<sup>-1</sup>.

EXAMPLE 4

10 Methyl 3-(Cyanomethyl)-2-[[[(4,6-dimethyl-  
2-pyrimidinyl)aminolcarbonyllaminol]sulfonyl]benzoate

A mixture of methyl 3-(cyanomethyl)-  
2-[[[(1,1-dimethylethyl)dimethylsilyl]amino]-  
sulfonyl]benzoate (0.34 g), O-phenyl-N-(4,6-dimethyl-  
15 2-pyrimidinyl)carbamate (0.24 g), and acetonitrile  
(2.5 mL) were stirred and cooled in an ice-acetone  
bath. A stock solution of 1 M tetrabutylammonium  
fluoride containing 8 mole % 1,8-diazabicyclo-  
[5.4.0]undec-7-ene (1.0 mL) was added and the  
20 reaction was stirred under a nitrogen atmosphere for  
45 minutes. The reaction was added to water (30 mL)  
and acidified to pH 5 with 1N hydrochloric acid. The  
resulting precipitate was filtered, washed with water  
and hexanes, and air dried to give 0.23 g of the  
25 title compound as a yellow solid, m.p. 182-185°C.

200 MHz NMR (DMSO) δ: 2.40 (s, 6H, CH<sub>3</sub>);  
3.79 (s, 3H, OCH<sub>3</sub>);  
4.67 (s, 2H, CH<sub>2</sub>CN);  
7.04 (s, 1H, pyrim. H);  
30 7.64 (m, 1H, arom.);  
7.82 (m, 2H, arom.);  
10.8 (s, 1H, NH); and  
13.7 (s, 1H, NH).

IR (mineral oil) 2240, 1740 cm<sup>-1</sup>.

35

EXAMPLE 5Methyl 3-Formyl-2-nitrobenzoate

Methyl 3-(dibromomethyl)-2-nitrobenzoate (62.6  
g), silver nitrate (109 g), 1,2-dimethoxyethane (500



15

mL), and water (400 mL) were heated at reflux overnight. The undissolved salts were filtered off  
5 and the filtrate was extracted with ethyl acetate. The extracts were washed with a brine solution, dried over magnesium sulfate, filtered, and concentrated to give a crude solid. Recrystallization from  
10 chlorobutane gave 20.9 g of the title compound as an orange solid.

90 MHz NMR (CDCl<sub>3</sub>) δ: 3.98 (s, 3H, OCH<sub>3</sub>);  
7.8 (m, 1H, arom.);  
8.3 (m, 2H, arom.); and  
10.0 (s, 1H, HCO).

15

EXAMPLE 6Methyl 3-(Difluoromethyl)-2-nitrobenzoate

To a solution of diethylaminosulfur trifluoride (11 mL) in dichloromethane (75 mL) stirred at -70°C under a nitrogen atmosphere, was added a  
20 dichloromethane solution of methyl 3-formyl-2-nitrobenzoate (5.9 g). The reaction was allowed to warm to room temperature. After 3 hours it was poured onto ice (300 mL) and extracted with  
dichloromethane. The extracts were washed with  
25 water, dried over magnesium sulfate, filtered, and concentrated to give a crude solid. Recrystallization from hexanes-chlorobutane gave 3.94 g of the title compound as a peach-colored solid, m.p. 66-69°C.

90 MHz NMR (CDCl<sub>3</sub>) δ: 3.94 (s, 3H, OCH<sub>3</sub>);  
30 6.87 (t, 1H, CHF<sub>2</sub>); and  
7.9 (m, 3H, arom.).

EXAMPLE 72-(Methoxycarbonyl)-6-(difluoromethyl)-benzenesulfonyl Chloride

35 To a stirred suspension of dry tetrahydrofuran (250 mL) and 26% potassium hydride in oil (6.17 g) was slowly added benzyl mercaptan (4.63 mL) at 0°C under a nitrogen atmosphere. After 15 minutes,

methyl 3-(difluor methyl)-2-nitrobenzoate (7.5 g) was added, and the resulting mixture was stirred at room temperature overnight. The reaction was contacted with aqueous sodium bicarbonate (50 mL) and extracted with ethyl acetate. The extracts were dried over magnesium sulfate, filtered, and concentrated to give 10.8 g of crude methyl 3-(difluoromethyl)-2-(phenylmethylthio)benzoate as a brown oil.

The crude thioether was stirred in a mixture of dichloromethane (400 mL) and 6N hydrochloric acid (28 mL) at 0°C, as 5% sodium hypochlorite (204 mL) was slowly added. The reaction mixture was stirred at 0°C for 2.5 hours. The dichloromethane layer was washed with water, dried over magnesium sulfate, filtered, concentrated, and triturated with hexanes to give 5.17 g of the title compound as a yellow solid.

90 MHz NMR (CDCl<sub>3</sub>) δ: 4.13 (s, 3H, OCH<sub>3</sub>); and 7.1 - 8.3 (m, 4H, CHF<sub>2</sub> and arom.).

#### EXAMPLE 8

Methyl 3-(Difluoromethyl)-2-[[[(1,1-dimethylethyl)dimethylsilyllaminolsulfonyl]benzoate

A suspension of 2-(methoxycarbonyl)-6-(difluoromethyl)benzenesulfonyl chloride (5.17 g), amino-t-butyldimethylsilane (4.9 g), and sodium bicarbonate (1.5 g) in dichloromethane (150 mL) was stirred at room temperature in a stoppered flask for 4 days. The organic phase was washed with water, dried over magnesium sulfate, filtered, concentrated, and chromatographed on silica gel eluting with 10 % ethyl acetate in hexanes to give 2.23 g of the title compound as a white solid.

90 MHz NMR (CDCl<sub>3</sub>) δ: 0.2 (s, 6H, SiCH<sub>3</sub>); 0.87 (s, 9H, C(CH<sub>3</sub>)<sub>3</sub>); 3.90 (s, 3H, OCH<sub>3</sub>);

17

5.88 (s, 1H, NH); and  
7.20 - 8.36 (m, 4H, CHF<sub>2</sub> and  
arom.).

5

EXAMPLE 9

Methyl 3-(Difluoromethyl)-2-[[[(4-methoxy-2-pyrimidinyl)aminolcarbonylaminol sulfonyl]benzoate

A mixture of methyl 3-(difluoromethyl)-2-  
10 [[[(1,1-dimethylethyl)dimethylsilyl]amino]sulfonyl]-  
benzoate (0.22 g), O-phenyl-N-(4-methoxy-  
2-pyrimidinyl)carbamate (0.15 g), and acetonitrile  
(2.0 mL) were stirred and cooled in an ice-acetone  
bath. A stock solution of 1 M tetrabutylammonium  
15 fluoride containing 8 mol % 1,8-diazabicyclo-  
[5.4.0]undec-7-ene (0.61 mL) was added, and the  
reaction was stirred under a nitrogen atmosphere for  
1 hour. The reaction was added to water (30 mL) and  
acidified to pH 5 with 1N hydrochloric acid. The  
20 resulting precipitate was filtered, washed with water  
and hexanes, and air dried to give 0.06 g of the  
title compound as a white solid, m.p. 178-180°C  
(decomposition).

200 MHz NMR (DMSO)  $\delta$ : 3.94 (s, 3H, OCH<sub>3</sub>);  
25 4.14 (s, 3H, OCH<sub>3</sub>);  
6.87 (d, 1H, pyrim. H);  
7.8 - 8.2 (m, 3H, arom.);  
8.43 (d, 1H, pyrim. H);  
11.46 (s, 1H, NH); and  
30 13.90 (s, 1H, NH).

IR (mineral oil) 1730, 1720 cm<sup>-1</sup>.

EXAMPLE 10

Methyl 3-(1-Hydroxyethyl)-2-nitrobenzoate

Titanium tetrachloride (6.25 mL) was added to  
35 dry diethylether (250 mL) at -78°C and was stirred  
under a nitrogen atmosphere. The resulting yellow  
suspension was allowed to warm to -50°C and was  
recooled to -78°C prior to adding 1.4 molar

methyllithium in diethylether (41.6 mL). The mixture was warmed to  $-30^{\circ}\text{C}$  and the resulting purple mixture was cooled to  $-60^{\circ}\text{C}$  and transferred by cannula into an adjacent flask containing methyl 3-formyl-2-nitrobenzoate (11.6 g) and dry tetrahydrofuran (250 mL). The resulting mixture was allowed to warm to room temperature and was stirred under a nitrogen atmosphere for about 18 hours. The reaction was poured onto a mixture of ice (100 mL) and 3N hydrochloric acid (100 mL) and extracted with ethyl acetate. The extracts were washed with saturated sodium bicarbonate and brine, dried over magnesium sulfate, filtered, and concentrated to a crude oil. Flash column chromatography on silica gel, eluting with dichloromethane gave 9.8 g of the title compound as a yellow oil,  $n_D$  1.5309.

90 MHz NMR ( $\text{CDCl}_3$ )  $\delta$ : 1.47 (d, 3H,  $\text{CH}_3$ );  
2.58 (s, 1H, OH);  
3.92 (s, 3H,  $\text{OCH}_3$ );  
4.93 (m, 1H, CH); and  
7.8 (m, 3H, arom.).

IR (neat) 3431, 1735  $\text{cm}^{-1}$ .

25

#### EXAMPLE 11

##### Methyl 3-(1-Fluoroethyl)-2-nitrobenzoate

To a solution of diethylamino sulfur trifluoride (18 mL), in dichloromethane (100 mL), stirred at  $-74^{\circ}\text{C}$  under a nitrogen atmosphere was added a dichloromethane solution of methyl 3-(1-hydroxyethyl)-2-nitrobenzoate (14.6 g). The reaction was allowed to warm to room temperature and was briefly warmed to  $35^{\circ}\text{C}$ . After a 4 hour reaction period, the reaction was poured onto ice and water (200 mL) and extracted with dichloromethane. The combined extracts were washed with a 1:1 mixture of 1N NaOH and saturated sodium bicarbonate, dried over magnesium sulfate, filtered, and concentrated to give

11.3 g of the title compound as a dark amber oil,  $n_D$  1.5151.

5 90 MHz NMR ( $CDCl_3$ )  $\delta$ : 1.67 (d of d, 3H,  $CH_3$ );  
3.93 (s, 3H,  $OCH_3$ );  
5.7 (d of q, 1H, CHF); and  
7.9 (m, 3H, arom.).

IR (neat) 1736  $cm^{-1}$ .

10

EXAMPLE 12

Methyl 3-(1-Fluoroethyl)-2-  
(phenylmethylthio)benzoate

Following the procedure described in Example 2,  
methyl 3-(1-fluoroethyl)-2-nitrobenzoate (15 g) was  
15 reacted to give 17.55 g of the unpurified title  
compound as a brown oil.

EXAMPLE 13

2-(Methoxycarbonyl)-6-(1-fluoroethyl)-  
benzene sulfonyl Chloride

20 Following the procedure described in Example 2,  
methyl 3-(1-fluoroethyl)-2-(phenylmethylthio)benzoate  
(8.8 g) was reacted and chromatographed on silica  
eluting with 10% ethyl acetate in hexanes to give 1.7  
g of the title compound as an orange oil.

25 90 MHz NMR ( $CDCl_3$ )  $\delta$ : 1.8 (d of d, 3H,  $CH_3$ );  
4.0 (s, 3H,  $OCH_3$ );  
6.6 (d of q, 1H, CHF); and  
7.8 (m, 3H, arom.).

EXAMPLE 14

30 Methyl 3-(1-Fluoroethyl)-2-[[[(1,1-dimethyl-  
ethyl)dimethylsilyl]aminol]sulfonyl]benzoate

Following the procedure described in Example 3,  
2-(methoxycarbonyl)-6-(1-fluoroethyl)benzene sulfonyl  
chloride (2.4 g) was reacted and chromatographed on  
35 silica luting with 15% thyl ac tat in hexan s t  
give 1.3 g of the title compound as an orang oil.

20

90 MHZ NMR (CDCl<sub>3</sub>) δ: 0.17 (s, 3H, SiCH<sub>3</sub>);  
0.24 (s, 3H, SiCH<sub>3</sub>);  
5 0.84 (s, 9H, C(CH<sub>3</sub>)<sub>3</sub>);  
1.57 (d of d, 3H, CH<sub>3</sub>);  
3.88 (s, 3H, OCH<sub>3</sub>);  
5.83 (s, 1H, NH);  
6.6 (d of q, 1H, CHF); and  
10 7.6 (m, 3H, arom.).

**EXAMPLE 15**

Methyl 3-(1-Fluoroethyl)-2-[[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]aminosulfonyl]benzoate

Following the procedure described in Example 4,  
15 methyl 3-(1-fluoroethyl)-2-[[[(1,1-dimethylethyl)-dimethylsilyl]amino]sulfonyl]benzoate (0.26 g) and O-phenyl-N-(4,6-dimethoxy-2-pyrimidinyl)carbamate (0.2 g) were reacted to give 0.16 g of the title compound as a pink solid, m.p. 155-158°C  
20 (decomposition).


200 MHz NMR (DMSO) δ: 1.64 (d of d, 3H, CH<sub>3</sub>);  
3.85 (s, 3H, CO<sub>2</sub>CH<sub>3</sub>);  
3.98 (s, 6H, OCH<sub>3</sub>);  
6.08 (s, 1H, pyrim. H);  
25 7.04 (d of q, 1H, CHF);  
7.72 (m, 1H, arom.);  
7.96 (m, 2H, arom.);  
10.85 (s, 1H, NH); and  
12.75 (s, 1H, NH).

30 IR (mineral oil) 3281, 1738, 1722 cm<sup>-1</sup>.

By applying the procedures of Equations 1 through 7 and Examples 1 through 15, the compounds in Tables 1 through 4 can be readily prepared by one skilled in the art.

35

TABLE 1

 $R^1$  is  $CH_3$ 

$R^2$	X	Y
$CH_2F$	$CH_3$	H
$CH_2F$	$CH_3$	$CH_3$
$CH_2F$	$CH_3$	$C_2H_5$
$CH_2F$	$CH_3$	$OCH_3$
$CH_2F$	$CH_3$	$OCHF_2$
$CH_2F$	$OCH_3$	H
$CH_2F$	$OCH_3$	Cl
$CH_2F$	$OCH_3$	$C_2H_5$
$CH_2F$	$OCH_3$	$OCH_3$
$CH_2F$	$OCH_3$	$OCHF_2$
$CHF_2$	$CH_3$	H
$CHF_2$	$CH_3$	$CH_3$
$CHF_2$	$CH_3$	$C_2H_5$
$CHF_2$	$CH_3$	$OCH_3$
$CHF_2$	$CH_3$	$OCH_2F$
$CHF_2$	$OCH_3$	H
$CHF_2$	$OCH_3$	Cl
$CHF_2$	$OCH_3$	$C_2H_5$

 $R^1$  is  $CH_2CH_3$ 

$R^2$	X	Y
$CH_2F$	$CH_3$	H
$CH_2F$	$CH_3$	$CH_3$
$CH_2F$	$CH_3$	$C_2H_5$
$CH_2F$	$CH_3$	$OCH_3$
$CH_2F$	$CH_3$	$OCHF_2$
$CH_2F$	$OCH_3$	H
$CH_2F$	$OCH_3$	Cl
$CH_2F$	$OCH_3$	$C_2H_5$
$CH_2F$	$OCH_3$	$OCH_3$
$CH_2F$	$OCH_3$	$OCHF_2$
$CHF_2$	$CH_3$	H
$CHF_2$	$CH_3$	$CH_3$
$CHF_2$	$CH_3$	$C_2H_5$
$CHF_2$	$CH_3$	$OCH_3$
$CHF_2$	$CH_3$	$OCH_2F$
$CHF_2$	$OCH_3$	H
$CHF_2$	$OCH_3$	Cl
$CHF_2$	$OCH_3$	$C_2H_5$

$R^1$  is  $CH_3$ 

$R^2$	X	Y
$CHF_2$	$OCH_3$	$OCH_3$
$CHF_2$	$OCH_3$	$OCHF_2$
$CH_2CN$	$CH_3$	H
$CH_2CN$	$CH_3$	$CH_3$
$CH_2CH$	$CH_3$	$C_2H_5$
$CH_2CN$	$CH_3$	$OCH_3$
$CH_2CN$	$CH_3$	$OCHF_2$
$CH_2CN$	$OCH_3$	H
$CH_2CN$	$OCH_3$	Cl
$CH_2CH$	$OCH_3$	$C_2H_5$
$CH_2CN$	$OCH_3$	$OCH_3$
$CH_2CN$	$OCH_3$	$OCHF_2$
$CHFCH_3$	$CH_3$	H
$CHFCH_3$	$CH_3$	$CH_3$
$CHFCH_3$	$CH_3$	$C_2H_5$
$CHFCH_3$	$CH_3$	$OCH_3$
$CHFCH_3$	$CH_3$	$OCHF_2$
$CHFCH_3$	$OCH_3$	H
$CHFCH_3$	$OCH_3$	Cl
$CHFCH_3$	$OCH_3$	$OCH_3$
$CHFCH_3$	$OCH_3$	$OCHF_2$

 $R^1$  is  $CH_2CH_2CH_3$ 

$R^2$	X	Y
$CH_2F$	$CH_3$	H
$CH_2F$	$CH_3$	$CH_3$
$CH_2F$	$CH_3$	$C_2H_5$
$CH_2F$	$CH_3$	$OCH_3$
$CH_2F$	$CH_3$	$OCHF_2$
$CH_2F$	$OCH_3$	H

 $R^1$  is  $CH_2CH_3$ 

$R^2$	X	Y
$CHF_2$	$OCH_3$	$OCH_3$
$CHF_2$	$OCH_3$	$OCHF_2$
$CH_2CN$	$CH_3$	H
$CH_2CN$	$CH_3$	$CH_3$
$CH_2CN$	$CH_3$	$C_2H_5$
$CH_2CN$	$CH_3$	$OCH_3$
$CH_2CN$	$CH_3$	$OCHF_2$
$CH_2CN$	$OCH_3$	H
$CH_2CN$	$OCH_3$	Cl
$CH_2CN$	$OCH_3$	$C_2H_5$
$CH_2CN$	$OCH_3$	$OCH_3$
$CH_2CN$	$OCH_3$	$OCHF_2$
$CHFCH_3$	$CH_3$	H
$CHFCH_3$	$CH_3$	$CH_3$
$CHFCH_3$	$CH_3$	$C_2H_5$
$CHFCH_3$	$CH_3$	$OCH_3$
$CHFCH_3$	$CH_3$	$OCHF_2$
$CHFCH_3$	$OCH_3$	H
$CHFCH_3$	$OCH_3$	Cl
$CHFCH_3$	$OCH_3$	$OCH_3$
$CHFCH_3$	$OCH_3$	$OCHF_2$

 $R^1$  is  $CH(CH_3)_2$ 

$R^2$	X	Y
$CH_2F$	$CH_3$	H
$CH_2F$	$CH_3$	$CH_3$
$CH_2F$	$CH_3$	$C_2H_5$
$CH_2F$	$CH_3$	$OCH_3$
$CH_2F$	$CH_3$	$OCHF_2$
$CH_2F$	$OCH_3$	H



$R^1$  is  $\text{CH}_2\text{CH}_2\text{CH}_3$ 

$R^2$	X	Y
$\text{CH}_2\text{F}$	$\text{OCH}_3$	Cl
$\text{CH}_2\text{F}$	$\text{OCH}_3$	$\text{C}_2\text{H}_5$
$\text{CH}_2\text{F}$	$\text{OCH}_3$	$\text{OCH}_3$
$\text{CH}_2\text{F}$	$\text{OCH}_3$	$\text{OCHF}_2$
$\text{CHF}_2$	$\text{CH}_3$	H
$\text{CHF}_2$	$\text{CH}_3$	$\text{CH}_3$
$\text{CHF}_2$	$\text{CH}_3$	$\text{C}_2\text{H}_5$
$\text{CHF}_2$	$\text{CH}_3$	$\text{OCH}_3$
$\text{CHF}_2$	$\text{CH}_3$	$\text{OCHF}_2$
$\text{CHF}_2$	$\text{OCH}_3$	H
$\text{CHF}_2$	$\text{OCH}_3$	Cl
$\text{CHF}_2$	$\text{OCH}_3$	$\text{C}_2\text{H}_5$
$\text{CHF}_2$	$\text{OCH}_3$	$\text{OCH}_3$
$\text{CHF}_2$	$\text{OCH}_3$	$\text{OCHF}_2$
$\text{CH}_2\text{CN}$	$\text{CH}_3$	H
$\text{CH}_2\text{CN}$	$\text{CH}_3$	$\text{CH}_3$
$\text{CH}_2\text{CN}$	$\text{CH}_3$	$\text{C}_2\text{H}_5$
$\text{CH}_2\text{CN}$	$\text{CH}_3$	$\text{OCH}_3$
$\text{CH}_2\text{CN}$	$\text{CH}_3$	$\text{OCHF}_2$
$\text{CH}_2\text{CN}$	$\text{OCH}_3$	H
$\text{CH}_2\text{CN}$	$\text{OCH}_3$	Cl
$\text{CH}_2\text{CN}$	$\text{OCH}_3$	$\text{C}_2\text{H}_5$
$\text{CH}_2\text{CN}$	$\text{OCH}_3$	$\text{OCH}_3$
$\text{CH}_2\text{CN}$	$\text{OCH}_3$	$\text{OCHF}_2$
$\text{CHFCH}_3$	$\text{CH}_3$	H
$\text{CHFCH}_3$	$\text{CH}_3$	$\text{CH}_3$
$\text{CHFCH}_3$	$\text{CH}_3$	$\text{C}_2\text{H}_5$
$\text{CHFCH}_3$	$\text{CH}_3$	$\text{OCH}_3$
$\text{CHFCH}_3$	$\text{CH}_3$	$\text{OCHF}_2$
$\text{CHFCH}_3$	$\text{OCH}_3$	H
$\text{CHFCH}_3$	$\text{OCH}_3$	Cl

 $R^1$  is  $\text{CH}(\text{CH}_3)_2$ 

$R^2$	X	Y
$\text{CH}_2\text{F}$	$\text{OCH}_3$	Cl
$\text{CH}_2\text{F}$	$\text{OCH}_3$	$\text{C}_2\text{H}_5$
$\text{CH}_2\text{F}$	$\text{OCH}_3$	$\text{OCH}_3$
$\text{CH}_2\text{F}$	$\text{OCH}_3$	$\text{OCHF}_2$
$\text{CHF}_2$	$\text{CH}_3$	H
$\text{CHF}_2$	$\text{CH}_3$	$\text{CH}_3$
$\text{CHF}_2$	$\text{CH}_3$	$\text{C}_2\text{H}_5$
$\text{CHF}_2$	$\text{CH}_3$	$\text{OCH}_3$
$\text{CHF}_2$	$\text{CH}_3$	$\text{OCH}_3$
$\text{CHF}_2$	$\text{OCH}_3$	H
$\text{CHF}_2$	$\text{OCH}_3$	Cl
$\text{CHF}_2$	$\text{OCH}_3$	$\text{C}_2\text{H}_5$
$\text{CHF}_2$	$\text{OCH}_3$	$\text{OCH}_3$
$\text{CHF}_2$	$\text{OCH}_3$	$\text{OCHF}_2$
$\text{CH}_2\text{CN}$	$\text{CH}_3$	H
$\text{CH}_2\text{CN}$	$\text{CH}_3$	$\text{CH}_3$
$\text{CH}_2\text{CN}$	$\text{CH}_3$	$\text{C}_2\text{H}_5$
$\text{CH}_2\text{CN}$	$\text{CH}_3$	$\text{OCH}_3$
$\text{CH}_2\text{CN}$	$\text{CH}_3$	$\text{OCHF}_2$
$\text{CH}_2\text{CN}$	$\text{OCH}_3$	H
$\text{CH}_2\text{CN}$	$\text{OCH}_3$	Cl
$\text{CH}_2\text{CN}$	$\text{OCH}_3$	$\text{C}_2\text{H}_5$
$\text{CH}_2\text{CN}$	$\text{OCH}_3$	$\text{OCH}_3$
$\text{CH}_2\text{CN}$	$\text{OCH}_3$	$\text{OCHF}_2$
$\text{CHFCH}_3$	$\text{CH}_3$	H
$\text{CHFCH}_3$	$\text{CH}_3$	$\text{CH}_3$
$\text{CHFCH}_3$	$\text{CH}_3$	$\text{C}_2\text{H}_5$
$\text{CHFCH}_3$	$\text{CH}_3$	$\text{OCH}_3$
$\text{CHFCH}_3$	$\text{CH}_3$	$\text{OCHF}_2$
$\text{CHFCH}_3$	$\text{OCH}_3$	H
$\text{CHFCH}_3$	$\text{OCH}_3$	Cl

$R^1$  is  $CH_2CH_2CH_3$

$R^2$	X	Y
$CHFCH_3$	$OCH_3$	$OCH_3$
$CHFCH_3$	$OCH_3$	$OCHF_2$

$R^1$  is  $CH_2CH=CH_2$

$R^2$	X	Y
$CH_2F$	$CH_3$	H
$CH_2F$	$CH_3$	$CH_3$
$CH_2F$	$CH_3$	$C_2H_5$
$CH_2F$	$CH_3$	$OCH_3$
$CH_2F$	$CH_3$	$OCHF_2$
$CH_2F$	$OCH_3$	H
$CH_2F$	$OCH_3$	Cl
$CH_2F$	$OCH_3$	$OCH_3$
$CH_2F$	$OCH_3$	$OCHF_2$
$CHF_2$	$CH_3$	H
$CHF_2$	$CH_3$	$CH_3$
$CHF_2$	$CH_3$	$C_2H_5$
$CHF_2$	$CH_3$	$OCH_3$
$CHF_2$	$CH_3$	$OCHF_2$
$CHF_2$	$OCH_3$	H
$CHF_2$	$OCH_3$	Cl
$CHF_2$	$OCH_3$	$OCH_3$
$CHF_2$	$OCH_3$	$OCHF_2$
$CH_2CN$	$CH_3$	H
$CH_2CN$	$CH_3$	$CH_3$
$CH_2CN$	$CH_3$	$C_2H_5$
$CH_2CN$	$CH_3$	$OCH_3$
$CH_2CN$	$CH_3$	$OCHF_2$
$CH_2CN$	$OCH_3$	H
$CH_2CN$	$OCH_3$	Cl

$R^1$  is  $CH(CH_3)_2$

$R^2$	X	Y
$CHFCH_3$	$OCH_3$	$OCH_3$
$CHFCH_3$	$OCH_3$	$OCHF_2$

$R^1$  is  $CH_2CH_2Cl$

$R^2$	X	Y
$CH_2F$	$CH_3$	H
$CH_2F$	$CH_3$	$CH_3$
$CH_2F$	$CH_3$	$OCH_3$
$CH_2F$	$CH_3$	$OCHF_2$
$CH_2F$	$OCH_3$	H
$CH_2F$	$OCH_3$	Cl
$CH_2F$	$OCH_3$	$C_2H_5$
$CH_2F$	$OCH_3$	$OCH_3$
$CH_2F$	$OCH_3$	$OCHF_2$
$CHF_2$	$CH_3$	H
$CHF_2$	$CH_3$	$CH_3$
$CHF_2$	$CH_3$	$OCH_3$
$CHF_2$	$CH_3$	$OCHF_2$
$CHF_2$	$OCH_3$	H
$CHF_2$	$OCH_3$	Cl
$CHF_2$	$OCH_3$	$C_2H_5$
$CHF_2$	$OCH_3$	$OCH_3$
$CHF_2$	$OCH_3$	$OCHF_2$
$CH_2CN$	$CH_3$	H
$CH_2CN$	$CH_3$	$CH_3$
$CH_2CN$	$CH_3$	$OCH_3$
$CH_2CN$	$CH_3$	$OCHF_2$
$CH_2CN$	$OCH_3$	H
$CH_2CN$	$OCH_3$	Cl
$CH_2CN$	$OCH_3$	$C_2H_5$

25

 $R^1$  is  $\text{CH}_2\text{CH}=\text{CH}_2$ 

$R^2$	X	Y
$\text{CH}_2\text{CN}$	$\text{OCH}_3$	$\text{OCH}_3$
$\text{CH}_2\text{CN}$	$\text{OCH}_3$	$\text{OCHF}_2$
$\text{CHFCH}_3$	$\text{CH}_3$	H
$\text{CHFCH}_3$	$\text{CH}_3$	$\text{CH}_3$
$\text{CHFCH}_3$	$\text{CH}_3$	$\text{C}_2\text{H}_5$
$\text{CHFCH}_3$	$\text{CH}_3$	$\text{OCH}_3$
$\text{CHFCH}_3$	$\text{CH}_3$	$\text{OCHF}_2$
$\text{CHFCH}_3$	$\text{OCH}_3$	H
$\text{CHFCH}_3$	$\text{OCH}_3$	Cl
$\text{CHFCH}_3$	$\text{OCH}_3$	$\text{OCH}_3$
$\text{CHFCH}_3$	$\text{OCH}_3$	$\text{OCHF}_2$

 $R^1$  is  $\text{CH}_2\text{C}\equiv\text{CH}$ 

$R^2$	X	Y
$\text{CH}_2\text{F}$	$\text{CH}_3$	H
$\text{CH}_2\text{F}$	$\text{CH}_3$	$\text{CH}_3$
$\text{CH}_2\text{F}$	$\text{CH}_3$	$\text{C}_2\text{H}_5$
$\text{CH}_2\text{F}$	$\text{CH}_3$	$\text{OCH}_3$
$\text{CH}_2\text{F}$	$\text{CH}_3$	$\text{OCHF}_2$
$\text{CH}_2\text{F}$	$\text{OCH}_3$	H
$\text{CH}_2\text{F}$	$\text{OCH}_3$	Cl
$\text{CH}_2\text{F}$	$\text{OCH}_3$	$\text{OCH}_3$
$\text{CH}_2\text{F}$	$\text{OCH}_3$	$\text{OCHF}_2$
$\text{CHF}_2$	$\text{CH}_3$	H
$\text{CHF}_2$	$\text{CH}_3$	$\text{CH}_3$
$\text{CHF}_2$	$\text{CH}_3$	$\text{C}_2\text{H}_5$
$\text{CHF}_2$	$\text{CH}_3$	$\text{OCH}_3$
$\text{CHF}_2$	$\text{CH}_3$	$\text{OCHF}_2$
$\text{CHF}_2$	$\text{OCH}_3$	H
$\text{CHF}_2$	$\text{OCH}_3$	Cl

 $R^1$  is  $\text{CH}_2\text{CH}_2\text{Cl}$ 

$R^2$	X	Y
$\text{CH}_2\text{CN}$	$\text{OCH}_3$	$\text{OCH}_3$
$\text{CH}_2\text{CN}$	$\text{OCH}_3$	$\text{OCHF}_2$
$\text{CHFCH}_3$	$\text{CH}_3$	$\text{CH}_3$
$\text{CHFCH}_3$	$\text{CH}_3$	$\text{CH}_3$
$\text{CHFCH}_3$	$\text{CH}_3$	$\text{C}_2\text{H}_5$
$\text{CHFCH}_3$	$\text{CH}_3$	$\text{OCH}_3$
$\text{CHFCH}_3$	$\text{CH}_3$	$\text{OCHF}_2$
$\text{CHFCH}_3$	$\text{OCH}_3$	H
$\text{CHFCH}_3$	$\text{OCH}_3$	Cl
$\text{CHFCH}_3$	$\text{OCH}_3$	$\text{OCH}_3$
$\text{CHFCH}_3$	$\text{OCH}_3$	$\text{OCHF}_2$

 $R^1$  is  $\text{CH}_2\text{CH}_2\text{OCH}_3$ 

$R^2$	X	Y
$\text{CH}_2\text{F}$	$\text{CH}_3$	H
$\text{CH}_2\text{F}$	$\text{CH}_3$	$\text{CH}_3$
$\text{CH}_2\text{F}$	$\text{CH}_3$	$\text{OCH}_3$
$\text{CH}_2\text{F}$	$\text{CH}_3$	$\text{OCHF}_2$
$\text{CH}_2\text{F}$	$\text{OCH}_3$	H
$\text{CH}_2\text{F}$	$\text{OCH}_3$	Cl
$\text{CH}_2\text{F}$	$\text{OCH}_3$	$\text{C}_2\text{H}_5$
$\text{CH}_2\text{F}$	$\text{OCH}_3$	$\text{OCH}_3$
$\text{CH}_2\text{F}$	$\text{OCH}_3$	$\text{OCHF}_2$
$\text{CHF}_2$	$\text{CH}_3$	H
$\text{CHF}_2$	$\text{CH}_3$	$\text{CH}_3$
$\text{CHF}_2$	$\text{CH}_3$	$\text{OCH}_3$
$\text{CHF}_2$	$\text{CH}_3$	$\text{OCHF}_2$
$\text{CHF}_2$	$\text{OCH}_3$	H
$\text{CHF}_2$	$\text{OCH}_3$	Cl
$\text{CHF}_2$	$\text{OCH}_3$	$\text{C}_2\text{H}_5$



$R^1$  is  $CH_2C\equiv CH$ 

$R^2$	X	Y
$CHF_2$	$OCH_3$	$OCH_3$
$CHF_2$	$OCH_3$	$OCHF_2$
$CH_2CN$	$CH_3$	H
$CH_2CN$	$CH_3$	$CH_3$
$CH_2CN$	$CH_3$	$C_2H_5$
$CH_2CN$	$CH_3$	$OCH_3$
$CH_2CN$	$CH_3$	$OCHF_2$
$CH_2CN$	$OCH_3$	H
$CH_2CN$	$OCH_3$	Cl
$CH_2CN$	$OCH_3$	$OCH_3$
$CH_2CN$	$OCH_3$	$OCHF_2$
$CHFCH_3$	$CH_3$	H
$CHFCH_3$	$CH_3$	$CH_3$
$CHFCH_3$	$CH_3$	$C_2H_5$
$CHFCH_3$	$CH_3$	$OCH_3$
$CHFCH_3$	$CH_3$	$OCHF_2$
$CHFCH_3$	$OCH_3$	H
$CHFCH_3$	$OCH_3$	Cl
$CHFCH_3$	$OCH_3$	$OCH_3$
$CHFCH_3$	$OCH_3$	$OCHF_2$

 $R^1$  is  $CH_2CH_2OCH_3$ 

$R^2$	X	Y
$CHF_2$	$OCH_3$	$OCH_3$
$CHF_2$	$OCH_3$	$OCHF_2$
$CH_2CN$	$CH_3$	H
$CH_2CN$	$CH_3$	$CH_3$
$CH_2CN$	$CH_3$	$OCH_3$
$CH_2CN$	$CH_3$	$OCHF_2$
$CH_2CN$	$OCH_3$	H
$CH_2CN$	$OCH_3$	Cl
$CH_2CN$	$OCH_3$	$C_2H_5$
$CH_2CN$	$OCH_3$	$OCH_3$
$CH_2CN$	$OCH_3$	$OCHF_2$
$CHFCH_3$	$CH_3$	H
$CHFCH_3$	$CH_3$	$CH_3$
$CHFCH_3$	$CH_3$	$C_2H_5$
$CHFCH_3$	$CH_3$	$OCH_3$
$CHFCH_3$	$CH_3$	$OCHF_2$
$CHFCH_3$	$OCH_3$	H
$CHFCH_3$	$OCH_3$	Cl
$CHFCH_3$	$OCH_3$	$OCH_3$
$CHFCH_3$	$OCH_3$	$OCHF_2$


27

**TABLE 3****TABLE 2**

$R^1$	$R^2$	$R^1$	$R^2$	$R^3$	$R^4$	$R^5$
CH <sub>3</sub>	CH <sub>2</sub> F	CH <sub>3</sub>	CH <sub>2</sub> F	CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>
CH <sub>3</sub>	CHF <sub>2</sub>	CH <sub>3</sub>	CHF <sub>2</sub>	CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>
CH <sub>3</sub>	CH <sub>2</sub> CN	CH <sub>3</sub>	CH <sub>2</sub> CN	CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>
CH <sub>3</sub>	CHFCH <sub>3</sub>	CH <sub>3</sub>	CHFCH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>
CH <sub>2</sub> CH <sub>3</sub>	CH <sub>2</sub> F	CH <sub>2</sub> CH <sub>3</sub>	CH <sub>2</sub> F	CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>
CH <sub>2</sub> CH <sub>3</sub>	CHF <sub>2</sub>	CH <sub>2</sub> CH <sub>3</sub>	CHF <sub>2</sub>	CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>
CH <sub>2</sub> CH <sub>3</sub>	CH <sub>2</sub> CN	CH <sub>2</sub> CH <sub>3</sub>	CH <sub>2</sub> CN	CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>
CH <sub>2</sub> CH <sub>3</sub>	CHFCH <sub>3</sub>	CH <sub>2</sub> CH <sub>3</sub>	CHFCH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>
CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	CH <sub>2</sub> F	CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	CH <sub>2</sub> F	CH <sub>3</sub>	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>
CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	CHF <sub>2</sub>	CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	CHF <sub>2</sub>	CH <sub>3</sub>	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>
CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	CH <sub>2</sub> CN	CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	CH <sub>2</sub> CN	CH <sub>3</sub>	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>
CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	CHFCH <sub>3</sub>	CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	CHFCH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>
CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>2</sub> F	CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>2</sub> F	CH <sub>3</sub>	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>
CH(CH <sub>3</sub> ) <sub>2</sub>	CHF <sub>2</sub>	CH(CH <sub>3</sub> ) <sub>2</sub>	CHF <sub>2</sub>	CH <sub>3</sub>	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>

R <sup>1</sup>	R <sup>2</sup>	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>
CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>2</sub> CN	CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>2</sub> CN	CH <sub>3</sub>	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>
CH(CH <sub>3</sub> ) <sub>2</sub>	CHFCH <sub>3</sub>	CH(CH <sub>3</sub> ) <sub>2</sub>	CHFCH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>
CH <sub>2</sub> CH=CH <sub>2</sub>	CH <sub>2</sub> F	CH <sub>2</sub> CH=CH <sub>2</sub>	CH <sub>2</sub> F	CH <sub>3</sub>	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>
CH <sub>2</sub> CH=CH <sub>2</sub>	CHF <sub>2</sub>	CH <sub>2</sub> CH=CH <sub>2</sub>	CHF <sub>2</sub>	CH <sub>3</sub>	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>
CH <sub>2</sub> CH=CH <sub>2</sub>	CH <sub>2</sub> CN	CH <sub>2</sub> CH=CH <sub>2</sub>	CH <sub>2</sub> CN	CH <sub>3</sub>	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>
CH <sub>2</sub> CH=CH <sub>2</sub>	CHFCH <sub>3</sub>	CH <sub>2</sub> CH=CH <sub>2</sub>	CHFCH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>
CH <sub>2</sub> C≡CH	CH <sub>2</sub> F	CH <sub>2</sub> C≡CH	CH <sub>2</sub> F	CH <sub>3</sub>	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>
CH <sub>2</sub> C≡CH	CHF <sub>2</sub>	CH <sub>2</sub> C≡CH	CHF <sub>2</sub>	CH <sub>3</sub>	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>
CH <sub>2</sub> C≡CH	CH <sub>2</sub> CN	CH <sub>2</sub> C≡CH	CH <sub>2</sub> CN	CH <sub>3</sub>	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>
CH <sub>2</sub> C≡CH	CHFCH <sub>3</sub>	CH <sub>2</sub> C≡CH	CHFCH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>
CH <sub>2</sub> CH <sub>2</sub> Cl	CH <sub>2</sub> F	CH <sub>2</sub> CH <sub>2</sub> Cl	CH <sub>2</sub> F	CH <sub>3</sub>	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>
CH <sub>2</sub> CH <sub>2</sub> Cl	CHF <sub>2</sub>	CH <sub>2</sub> CH <sub>2</sub> Cl	CHF <sub>2</sub>	CH <sub>3</sub>	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>
CH <sub>2</sub> CH <sub>2</sub> Cl	CH <sub>2</sub> CN	CH <sub>2</sub> CH <sub>2</sub> Cl	CH <sub>2</sub> CN	CH <sub>3</sub>	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>
CH <sub>2</sub> CH <sub>2</sub> Cl	CHFCH <sub>3</sub>	CH <sub>2</sub> CH <sub>2</sub> Cl	CHFCH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>
CH <sub>2</sub> CH <sub>2</sub> OCH <sub>3</sub>	CH <sub>2</sub> F	CH <sub>2</sub> CH <sub>2</sub> OCH <sub>3</sub>	CH <sub>2</sub> F	CH <sub>3</sub>	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>
CH <sub>2</sub> CH <sub>2</sub> OCH <sub>3</sub>	CHF <sub>2</sub>	CH <sub>2</sub> CH <sub>2</sub> OCH <sub>3</sub>	CHF <sub>2</sub>	CH <sub>3</sub>	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>
CH <sub>2</sub> CH <sub>2</sub> OCH <sub>3</sub>	CH <sub>2</sub> CN	CH <sub>2</sub> CH <sub>2</sub> OCH <sub>3</sub>	CH <sub>2</sub> CN	CH <sub>3</sub>	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>
CH <sub>2</sub> CH <sub>2</sub> OCH <sub>3</sub>	CHFCH <sub>3</sub>	CH <sub>2</sub> CH <sub>2</sub> OCH <sub>3</sub>	CHFCH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>

TABLE 4



R <sup>2</sup>	X	Y	R <sup>2</sup>	X	Y
CH <sub>2</sub> F	CH <sub>3</sub>	H	CH <sub>2</sub> CN	CH <sub>3</sub>	H
CH <sub>2</sub> F	CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>2</sub> CN	CH <sub>3</sub>	CH <sub>3</sub>
CH <sub>2</sub> F	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> CN	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>
CH <sub>2</sub> F	CH <sub>3</sub>	OCH <sub>3</sub>	CH <sub>2</sub> CN	CH <sub>3</sub>	OCH <sub>3</sub>
CH <sub>2</sub> F	CH <sub>3</sub>	OCHF <sub>2</sub>	CH <sub>2</sub> CN	CH <sub>3</sub>	OCHF <sub>2</sub>
CH <sub>2</sub> F	OCH <sub>3</sub>	H	CH <sub>2</sub> CN	OCH <sub>3</sub>	H
CH <sub>2</sub> F	OCH <sub>3</sub>	Cl	CH <sub>2</sub> CN	OCH <sub>3</sub>	Cl
CH <sub>2</sub> F	OCH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> CN	OCH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>
CH <sub>2</sub> F	OCH <sub>3</sub>	OCH <sub>3</sub>	CH <sub>2</sub> CN	OCH <sub>3</sub>	OCH <sub>3</sub>
CH <sub>2</sub> F	OCH <sub>3</sub>	OCHF <sub>2</sub>	CH <sub>2</sub> CN	OCH <sub>3</sub>	OCHF <sub>2</sub>
CHF <sub>2</sub>	CH <sub>3</sub>	H	CHFCH <sub>3</sub>	CH <sub>3</sub>	H
CHF <sub>2</sub>	CH <sub>3</sub>	CH <sub>3</sub>	CHFCH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>
CHF <sub>2</sub>	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	CHFCH <sub>3</sub>	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>
CHF <sub>2</sub>	CH <sub>3</sub>	OCH <sub>3</sub>	CHFCH <sub>3</sub>	CH <sub>3</sub>	OCH <sub>3</sub>
CHF <sub>2</sub>	CH <sub>3</sub>	OCHF <sub>2</sub>	CHFCH <sub>3</sub>	CH <sub>3</sub>	OCHF <sub>2</sub>
CHF <sub>2</sub>	OCH <sub>3</sub>	H	CHFCH <sub>3</sub>	OCH <sub>3</sub>	H
CHF <sub>2</sub>	OCH <sub>3</sub>	Cl	CHFCH <sub>3</sub>	OCH <sub>3</sub>	Cl
CHF <sub>2</sub>	OCH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	CHFCH <sub>3</sub>	OCH <sub>3</sub>	OCH <sub>3</sub>
CHF <sub>2</sub>	OCH <sub>3</sub>	OCH <sub>3</sub>	CHFCH <sub>3</sub>	OCH <sub>3</sub>	OCHF <sub>2</sub>
CHF <sub>2</sub>	OCH <sub>3</sub>	OCHF <sub>2</sub>			

**Formulations**

Useful formulations of the compounds of Formula I can be prepared in conventional ways. They include dusts, granules, pellets, solutions, suspensions, emulsions, wettable powders, emulsifiable concentrates and the like. Many of these may be applied directly. Sprayable formulations can be extended in suitable media and used at spray volumes of from a few liters to several hundred liters per hectare. High strength compositions are primarily used as intermediates for further formulation. The formulations, broadly, contain about 0.1% to 99% by weight of active ingredient(s) and at least one of (a) about 0.1% to 20% surfactant(s) and (b) about 1% to 99.9% solid or liquid inert diluent(s). More specifically, they will contain these ingredients in the following approximate proportions:

		<u>Weight Percent*</u>		
		<u>Active</u>		
		<u>Ingredient</u>	<u>Diluent(s)</u>	<u>Surfactant(s)</u>
	Wettable Powders	20-90	0-74	1-10
25	Oil Suspensions, Emulsions, Solutions, (including Emulsifiable Concentrates)	3-50	40-95	0-15
	Aqueous Suspension	10-50	40-84	1-20
30	Dusts	1-25	70-99	0-5
	Granules and Pellets	0.1-95	5-99.9	0-15
	High Strength Compositions	90-99	0-10	0-2

35 \* Active ingredient plus at least one of a Surfactant or a Diluent equals 100 weight percent.



Lower or high levels of active ingredient can, of course, be present depending on the intended use and the physical properties of the compound. Higher ratios of surfactant to active ingredient are sometimes desirable, and are achieved by incorporation into the formulation or by tank mixing.

Typical solid diluents are described in Watkins, et al., "Handbook of Insecticide Dust Diluents and Carriers", 2nd Ed., Dorland Books, Caldwell, New Jersey, but other solids, either mined or manufactured, may be used. The more absorptive diluents are preferred for wettable powders and the denser ones for dusts. Typical liquid diluents and solvents are described in Marsden, "Solvents Guide," 2nd Ed., Interscience, New York, 1950. Solubility under 0.1% is preferred for suspension concentrates; solution concentrates are preferably stable against phase separation at 0°C. "McCutcheon's Detergents and Emulsifiers Annual", MC Publishing Corp., Ridgewood, New Jersey, as well as Sisely and Wood, "Encyclopedia of Surface Active Agents", Chemical Publishing Co., Inc., New York, 1964, list surfactants and recommended uses. All formulations can contain minor amounts of additives to reduce foaming, caking, corrosion, microbiological growth, etc.

The methods of making such compositions are well known. Solutions are prepared by simply mixing the ingredients. Fine solid compositions are made by blending and, usually, grinding as in a hammer or fluid energy mill. Suspensions are prepared by wet milling (see, for example, Littler, U.S. Patent 3,060,084). Granules and pellets may be made by spraying the active material upon preformed granular carriers or by agglomeration technique. S. J. E. Browning, "Agglomeration", Chemical Engineering, December 4, 1967, pp. 147ff. and "Perry's Chemical

Engineer's Handbook", 5th Ed., McGraw-Hill, New York, 1963, pp. 8-57ff.

5 For further information regarding the art of formulation, see for example:

H. M. Loux, U.S. Patent 3,235,361, February 15, 1966, Col. 6, line 16 through Col. 7, line 19 and Examples 10 through 41;

10 R. W. Luckenbaugh, U.S. Patent 3,309,192, March 14, 1967, Col. 5, line 43 through Col. 7, line 62 and Examples 8, 12, 15, 39, 41, 52, 53, 58, 132, 138-140, 162-164, 166, 167 and 169-182;

15 H. Gysin and E. Knusli, U.S. Patent 2,891,855, June 23, 1959, Col. 3, line 66 through Col. 5, line 17 and Examples 1-4; G. C. Klingman, "Weed Control as a Science", John Wiley and Sons, Inc., New York, 1961, pp. 81-96; and

20 J. D. Fryer and S. A. Evans, "Weed Control Handbook", 5th Ed., Blackwell Scientific Publications, Oxford, 1968, pp. 101-103.

In the following examples, all parts are by weight unless otherwise indicated.

Example A

High Strength Concentrate

Methyl 3-(cyanomethyl)-2-[[[(4,6-dimethyl-2-pyrimidinyl)amino]carbonyl]amino]-

30 sulfonyl]benzoate 99%  
trimethylnonyl polyethylene glycol ether 1%

The surfactant is sprayed upon the active ingredient in a blender and the mixture sifted through a U. S. S. No. 40 sieve (0.42 mm openings) prior to packaging. The concentrate may be formulated further for practical use.

Example BWettable Powder

5	Methyl 2-[[[(4-chloro-6-methoxy-2-pyrimidinyl)-amino]carbonyl]amino]sulfonyl]-3-(cyanomethyl)-benzoate	65%
	dodecylphenol polyethylene glycol ether	2%
	sodium ligninsulfonate	4%
10	sodium silicoaluminate	6%
	montmorillonite (calcined)	23%

The ingredients are thoroughly blended. The liquid surfactant is added by spraying upon the solid ingredients in the blender. After grinding in a hammer mill to produce particles essentially all below 100 microns, the material is reblended and sifted through a U.S.S. No. 50 sieve (0.3 mm opening) and packaged.

Example C20 Aqueous Suspension

	Methyl 3-(cyanomethyl)-2-[[[(4-(difluoromethoxy)-6-methoxy-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]benzoate	50.0%
	polyacrylic acid thickener	0.3%
25	dodecylphenol polyethylene glycol ether	0.5%
	disodium phosphate	1%
	monosodium phosphate	0.5%
	polyvinyl alcohol	1.0%
	water	56.7%

30 The ingredients are blended and ground together in a sand mill to produce particles essentially all under 5 microns in size.

Example DOil Suspension

35	Methyl 3-(difluoromethyl)-2-[[[(4-methoxy-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]-benzoate	35%
----	---	-----

- blend of polyalcohol carboxylic esters  
and oil soluble petroleum sulfonates 6%
- 5 xylene 59%

The ingredients are combined and ground together in a sand mill to produce particles essentially all below 3 microns. The product can be used directly, extended with oils, or emulsified in water.

10 Example E

Oil Suspension

- Methyl 3-(fluoromethyl)-2-[[[(4-methoxy-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]-benzoate 25%
- 15 polyoxyethylene sorbitol hexaoleate 5%
- highly aliphatic hydrocarbon oil 70%

The ingredients are ground together in a sand mill until the solid particles have been reduced to under about 5 microns. The resulting thick suspension may be applied directly, but preferably after being extended with oils or emulsified in water.

Example F

Aqueous Suspension

- Methyl 2-[[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]-3-(fluoromethyl)-benzoate 25%
- 25 hydrated attapulgite 3%
- crude calcium ligninsulfonate 10%
- sodium dihydrogen phosphate 0.5%
- 30 water 61.5%

The ingredients are ground together in a ball or roller mill until the solid particles have been reduced to diameters under 10 microns.

Example G

35 Wettable Powder

- Ethyl 2-[[[(4-chloro-6-methoxy-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]-3-(fluoromethyl)-benzoate 40.0%

## 35

	dioctyl sodium sulfosuccinate	1.5%
	sodium ligninsulfonate	3%
5	low viscosity methyl cellulose	1.5%
	attapulgate	54%

The ingredients are thoroughly blended, passed through an air mill, to produce an average particle size under 15 microns, reblended, and sifted through a U.S.S. No. 50 sieve (0.3 mm opening) before packaging.

All compounds of the invention may be formulated in the same manner.

Example HGranule

15	wettable powder of Example G	15%
	gypsum	69%
	potassium sulfate	16%

The ingredients are blended in a rotating mixer and water sprayed on to accomplish granulation. When most of the material has reached the desired range of 1.0 to 0.42 cm (U.S.S.#18 to 40 sieves), the granules are removed, dried, and screened. Oversized material is crushed to produce additional material in the desired range. These granules contain % active ingredient.

Example IWettable Powder

	Methyl 2-[[[(4,6-dimethoxy-2-pyrimidinyl)amino]-carbonyl]amino]sulfonyl]-3-(fluoromethyl)-	
30	benzoate	50%
	sodium alkyl naphthalenesulfonate	2%
	low viscosity methyl cellulose	2%
	diatomaceous earth	46%

The ingredients are blended, coarsely hammer-milled and then air milled to produce particles of active essentially all below 10 microns in diameter. The product is reblended before packaging.

Example JExtruded Pellet

5	Methyl 3-(fluoromethyl)-2-[[[(4-methoxy-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]-benzoate	25%
	anhydrous sodium sulfate	10%
	crude calcium ligninsulfonate	5%
10	sodium alkyl naphthalenesulfonate	1%
	calcium/magnesium bentonite	59%

The ingredients are blended, hammer-milled and then moistened with about 12% water. The mixture is extruded as cylinders about 3 mm diameter which are cut to produce pellets about 3 mm long. These may be used directly after drying, or the dried pellets may be crushed to pass a U.S.S. No. 20 sieve (0.84 mm openings). The granules held on a U.S.S. No. 40 sieve (0.42 mm openings) may be packaged for use and the fines recycled.

Example KWettable Powder

25	Methyl 3-(difluoromethyl)-2-[[[(4-methoxy-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]-benzoate	80%
	sodium alkyl naphthalenesulfonate	2%
	sodium ligninsulfonate	2%
	synthetic amorphous silica	3%
	kaolinite	13%

The ingredients are blended and then ground in a hammermill to produce particles with an average particle size less than 25 microns in diameter. The material is reblended and sifted through a U.S.S. No. 50 sieve (0.3 mm opening) before being packaged.

### Example 1

## High Strength Concentrate

- |   |  |       |
|---|--|-------|
| 5 | Methyl 3-(cyanomethyl)-2-[[[(4-(difluoromethoxy)-6-methoxy-2-pyrimidinyl]amino]carbonyl]amino]-sulfonyl]benzoate | 98.5% |
|   | silica aerogel   | 0.5%  |
|   | synthetic amorphous fine silica  | 1.0%  |
- 10           The ingredients are blended and ground in a hammer mill to produce a high strength concentrate essentially all passing a U.S.S. No. 50 sieve (0.3 mm openings). This material may then be formulated in a variety of ways.

### Example M

### Solution

- |    |  |     |
|----|--|-----|
|    | Methyl 3-(cyanomethyl)-2-[[[(4,6-dimethyl-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]-benzoate, sodium salt | 5%  |
| 20 | water  | 95% |

The salt is added directly to the water with stirring to produce the solution, which may then be packaged for use.

**25      UTILITY**

Test results indicate that compounds of this invention are active postemergence and preemergence herbicides. Many compounds of this invention are useful for the control of selected grass and broad-  
30 leaf weeds with tolerance to important agronomic crops such as barley (Hordeum vulgare), corn (Zea mays), rice (Oryza sativa), and wheat (Triticum aestivum). Grass weeds controlled include, but are not limited to, barnyardgrass (Echinochloa crus-galli), black-  
35 grass (Alopecurus myosuroides), Bromus spp., fxtail (Setaria spp.), johnsongrass (Sorghum halepense), panicum (Panicum spp.), and wild oat (Avena fatua). Broadleaf weeds controlled include, but are not

limited to, cocklebur (Xanthium pensylvanicum),  
jimsonweed (Datura stramonium), lambsquarters  
5 (Chenopodium album), morningglory (Ipomoea spp.),  
pigweed (Amaranthus spp.), Polygonum spp., sicklepod  
(Cassia obtusifolia), and velvetleaf (Abutilon  
theophrasti). Many compounds in this invention also  
control nutsedge (Cyperus spp.). Several compounds  
10 from this invention are particularly useful for weed  
control in cereal crops such as barley and wheat. A  
select few of these compounds are particularly useful  
for weed control in corn.

Several compounds in this invention have utility  
15 in non-crop areas where selected or complete control  
of plants or weeds is desired, such as around storage  
tanks, parking lots, billboards, highways, and  
railroad structures. These compounds are also useful  
in fallow areas of crop production such as in wheat  
20 and barley and in plantation crops such as palm,  
banana, citrus, rubber, etc. Alternatively, these  
compounds may be useful to modify plant growth or as  
citrus harvest aid abscission agents.

Rates of application for compounds of this  
25 invention are determined by a number of factors.  
These factors include: formulation selected, method  
of application, amount of vegetation present, growing  
conditions, etc. In general terms, the subject  
compounds should be applied at rates from 0.001 to 20  
30 kg/ha with a preferred rate range of from 0.004 to  
0.25 kg/ha. One skilled in the art can easily  
determine rates needed for the desired level of weed  
control.

Compounds of this invention may be used alone or  
35 in combination with other commercial herbicides,  
insecticides, or fungicides. The following list



exemplifies some of the herbicides suitable for use in mixtures. A combination of compounds from this invention with one or more of the following herbicides may be particularly useful for weed control.

	<u>Common Name</u>	<u>Chemical Name</u>
10	acetochlor	2-chloro-N-(ethoxymethyl)-N-(2-ethyl-6-methylphenyl)acetamide
	acifluorfen	5-[2-chloro-4-(trifluoromethyl)-phenoxy]-2-nitrobenzoic acid
	alachlor	2-chloro-N-(2,6-diethylphenyl)-N-(methoxymethyl)acetamide
15	anilofos	S-4-chloro-N-isopropylcarbaniloyl-methyl-O,O-dimethyl phosphorodithioate
	ametryn	N-ethyl-N'-(1-methylethyl)-6-(methylthio)-1,3,5-triazine-2,4-diamine
20	amitrole	1H-1,2,4-triazol-3-amine
	AMS	ammonium sulfamate
	asulam	methyl [(4-aminophenyl)sulfonyl]-carbamate
25	atrazine	6-chloro-N-ethyl-N'-(1-methylethyl)-1,3,5-triazine-2,4-diamine
	barban	4-chloro-2-butynyl 3-chlorocarbamat
	benefin	N-butyl-N-ethyl-2,6-dinitro-4-(trifluoromethyl)benzenamine
30	bensulfuron methyl	2-[[[[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]-amino]sulfonyl]methyl]benzoic acid, methyl ester
	bensulide	O,O-bis(1-methylethyl) S-[2-[(phenylsulfonyl)amino]-thyl]phosphorodithioat
35	b ntazon	3-(1-methylethyl)-(1H)-2,1,3-benzothiadiazin-4(3H)-on , 2,2-dioxide

	<u>Common Name</u>	<u>Chemical Name</u>
5	benzofluor	N-[4-(ethylthio)-2-(trifluoromethyl)phenyl]methanesulfonamide
	benzoylprop	N-benzoyl-N-(3,4-dichlorophenyl)-DL-alanine
10	bifenox	methyl 5-(2,4-dichlorophenoxy)-2-nitrobenzoate
	bromacil	5-bromo-6-methyl-3-(1-methylpropyl)-2,4(1H,3H)pyrimidinedione
	bromoxynil	3,5-dibromo-4-hydroxybenzonitrile
15	butachlor	N-(butoxymethyl)-2-chloro-N-(2,6-diethylphenyl)acetamide
	buthidazole	3-[5-(1,1-dimethylethyl)-1,3,4-thiadiazol-2-yl]-4-hydroxy-1-methyl-2-imidazolidinone
20	butralin	4-(1,1-dimethylethyl)-N-(1-methylpropyl)-2,6-dinitrobenzenamine
	butylate	S-ethyl bis(2-methylpropyl)-carbamothioate
	cacodylic acid	dimethyl arsinic oxide
25	CDAA	2-chloro-N,N-di-2-propenylacetamide
	CDEC	2-chloroallyl diethyldithiocarbamate
	CGA 142,464	3-(4,6-dimethoxy-1,3,5-triazin-2-yl)-1-[2-(2-methoxyethoxy)-phenylsulfonyl]-urea
30	chloramben	3-amino-2,5-dichlorobenzoic acid
	chlorbromuron	3-(4-bromo-3-chlorophenyl)-1-methoxy-1-methylurea
35	chlorimuron ethyl	2-[[[(4-chloro-6-methoxy-2-pyrimidinyl)amino]carbonyl]-amino]sulfonyl]benzoic acid, ethyl ester
	chlormethoxy-nil	2,4-dichlorophenyl 4-nitro-3-methoxyphenylether

	<u>Common Name</u>	<u>Chemical Name</u>
5	chlornitrofen	2,4,6-trichlorophenyl-4-nitrophenyl ether
	chloroxuron	N'-[4-(4-chlorophenoxy)phenyl]-N,N-dimethylurea
	chlorpropham	1-methylethyl 3-chlorophenylcarbamate
10	chlorsulfuron	2-chloro-N-[[4-methoxy-6-methyl-1,3,5-triazin-2-yl]amino]carbonyl]benzenesulfonamide
	chlortoluron	N'-(3-chloro-4-methylphenyl)-N,N-dimethylurea
15	cinmethylin	exo-1-methyl-4-(1-methylethyl)-2-[(2-methylphenyl)methoxy]-7-oxabicyclo[2.2.1]heptane
	clethodim	(E,E)-(±)-2-[1-[[3-chloro-2-propenyl]oxy]imino]propyl]-5-[2-(ethylthio)propyl]-3-hydroxy-2-cyclohexen-1-one
20	clomazone	2-[(2-chlorophenyl)methyl]-4,4-dimethyl-3-isoxazolidinone
	cloproxydim	(E,E)-2-[1-[[3-chloro-2-propenyl]oxy]imino]butyl]-5-[2-(ethylthio)propyl]-3-hydroxy-2-cyclohexen-1-one
25	clopyralid	3,6-dichloro-2-pyridinecarboxylic acid
	CMA	calcium salt of MAA
	cyanazine	2-[[4-chloro-6-(ethylamino)-1,3,5-triazin-2-yl]amino]-2-methylpropanenitril
30	cycloate	S-ethyl cyclohexylethylcarbamothioate
	cycluron	3-cyclooctyl-1,1-dimethylurea
	cyperquat	1-methyl-4-phenylpyridinium
	cyprazine	2-chloro-4-(cyclopropylamino)-6-(isopropylamino)-s-triazine
35		

	<u>Common Name</u>	<u>Chemical Name</u>
5	cyprazole	N-[5-(2-chloro-1,1-dimethylethyl)-1,3,4-thiadiazol-2-yl]cyclopropanecarboxamide
	cypromid	3',4'-dichlorocyclopropanecarboxanilide
	dalapon	2,2-dichloropropanoic acid
10	dazomet	tetrahydro-3,5-dimethyl-2H-1,3,5-thiadiazine-2-thione
	DCPA	dimethyl 2,3,5,6-tetrachloro-1,4-benzenedicarboxylate
15	desmediphan	ethyl [3-[[[(phenylamino)carbonyl]oxy]-phenyl]carbamate
	desmetryn	2-(isopropylamino)-4-(methylamino)-6-(methylthio)-s-triazine
	diallate	S-(2,3-dichloro-2-propenyl)bis(1-methylethyl)carbamoethioate
20	dicamba	3,6-dichloro-2-methoxybenzoic acid
	dichlobenil	2,6-dichlorobenzonitrile
	dichlorprop	(±)-2-(2,4-dichlorophenoxy)propanoic acid
25	dichlofop	(±)-2-[4-(2,4-dichlorophenoxy)phenoxy]propanoic acid, methyl ester
	diethatyl	N-(chloroacetyl)-N-(2,6-diethylphenyl)-glycine
	difenzoquat	1,2-dimethyl-3,5-diphenyl-1H-pyrazolium
30	dimepiperate	S-1-methyl-1-phenylethylpiperidine-1-carbothioate
	dinitramine	N <sup>3</sup> ,N <sup>3</sup> -diethyl-2,4-dinitro-6-(trifluoromethyl)-1,3-benzenediamine
35	dinoseb	2-(1-methylpropyl)-4,6-dinitrophenol
	diphenamid	N,N-dimethyl-α-phenylbenz n acetamide

	<u>Common Name</u>	<u>Chemical Name</u>
5	dipropetryn	6-(ethylthio)-N,N'-bis(1-methylethyl)-1,3,5-triazine-2,4-diamine
	diquat	6,7-dihydrodipyrido[1,2-a:2',1'-c]-pyrazinedium ion
	diuron	N'-(3,4-dichlorophenyl)-N,N-dimethylurea
10	DNOC	2-methyl-4,6-dinitrophenol
	DPX-M6316	3-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)amino]carbonyl]amino]sulfonyl]-2-thiophenecarboxylic acid, methyl ester
15	DSMA	disodium salt of MAA
	dymron	N-(4-methylphenyl)-N'-(1-methyl-1-phenylethyl)urea
	endothall	7-oxabicyclo[2.2.1]heptane-2,3-dicarboxylic acid
20	EPTC	S-ethyl dipropylcarbamoithioate
	esprocarb (SC2957)	S-benzyl-N-ethyl-N-(1,2-dimethyl)-propylthiolcarbamate
	ethalfluralin	N-ethyl-N-(2-methyl-2-propenyl)-2,6-dinitro-4-(trifluoromethyl)-benzenamine
25	ethofumesate	(±)-2-ethoxy-2,3-dihydro-3,3-dimethyl-5-benzofuranyl methanesulfonate
	fenac	2,3,6-trichlorobenzeneacetic acid
30	fenoxaprop	(±)-2-[4-[(6-chloro-2-benzoxazolyl)oxy]-phenoxy]propanoic acid
	fenuron	N,N-dimethyl-N'-phenylurea
	fenuron TCA	Salt of fenuron and TCA
35	flamprop	N-benzoyl-N-(3-chloro-4-fluorophenyl)-DL-alanine

	<u>Common Name</u>	<u>Chemical Name</u>
5	fluazifop	(±)-2-[4-[[5-(trifluoromethyl)-2-pyridinyl]oxy]phenoxy]propanoic acid
	fluazifop-P	(R)-2-[4-[[5-(trifluoromethyl)-2-pyridinyl]oxy]phenoxy]propanoic acid
10	fluchloralin	N-(2-chloroethyl)-2,6-dinitro-N-propyl-4-(trifluoromethyl)benzenamine
	fluometuron	N,N-dimethyl-N'-[3-(trifluoromethyl)-phenyl]urea
	fluorochloridone	3-chloro-4-(chloromethyl)-1-[3-(trifluoromethyl)phenyl]-2-pyrrolidinone
15	fluorodifen	p-nitrophenyl α,α,α-trifluoro-2-nitro-p-tolyl ether
	fluoroglycofen	carboxymethyl 5-[2-chloro-4-(trifluoromethyl)phenoxy]-2-nitrobenzoate
20	fluridone	1-methyl-3-phenyl-5-[3-(trifluoromethyl)phenyl]-4(1H)-pyridinone
	fomesafen	5-[2-chloro-4-(trifluoromethyl)phenoxy]-N-(methylsulfonyl)-2-nitrobenzamide
	fosamine	ethyl hydrogen (aminocarbonyl)-phosphate
25	glyphosate	N-(phosphonomethyl)glycine
	haloxyfop	2-[4-[[3-chloro-5-(trifluoromethyl)-2-pyridinyl]oxy]phenoxy]propanoic acid
	hexaflurate	potassium hexafluoroarsenate
30	hexazinone	3-cyclohexyl-6-(dimethylamino)-1-methyl-1,3,5-triazine-2,4(1H,3H)-dione
	imazamethabenz	6-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-m-toluic acid, methyl ester and 6-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-p-toluic acid, methyl ester
35		

	<u>Common Name</u>	<u>Chemical Name</u>
5	imazapyr	(±)-2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-pyridinecarboxylic acid
	imazaquin	2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-quinolinecarboxylic acid
10	imazethapyr	(±)-2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-ethyl-3-pyridinecarboxylic acid
	ioxynil	4-hydroxy-3,5-diiodobenzonitrile
15	isopropalin'	4-(1-methylethyl)-2,6-dinitro-N,N-dipropylbenzenamine
	isoproturon	N-(4-isopropylphenyl)-N',N'-dimethylurea
	isouron	N'-[5-(1,1-dimethylethyl)-3-isoxazolyl]-N,N-dimethylurea
20	isoxaben	N-[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide
	karbutilate	3-[(dimethylamino)carbonylamino]-phenyl-(1,1-dimethylethyl)carbamate
25	lactofen	(±)-2-ethoxy-1-methyl-2-oxoethyl 5-[2-chloro-4-(trifluoromethyl)phenoxy]-2-nitrobenzoate
	lenacil	3-cyclohexyl-6,7-dihydro-1H-cyclopentapyrimidine-2,4(3H,5H)-dione
	linuron	N'-(3,4-dichlorophenyl)-N-methoxy-N-methylurea
30	MAA	methylarsonic acid
	MAMA	monoammonium salt of MAA
	MCPA	(4-chloro-2-methylphenoxy)acetic acid
35	MCPB	4-(4-chloro-2-methylphenoxy)butanoic acid

	<u>Common Name</u>	<u>Chemical Name</u>
5	MON 7200	S,S-dimethyl-2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-3,5-pyridinedicarbothionate
	mecoprop	(±)-2-(4-chloro-2-methylphenoxy)-propanoic acid
10	mefenacet	2-(2-benzothiazolyloxy-N-methyl-N-phenylacetamide
	mefluidide	N-[2,4-dimethyl-5-[[[(trifluoromethyl)-sulfonyl]amino]phenyl]acetamide
15	methal-propalin	N-(2-methyl-2-propenyl)-2,6-dinitro-N-propyl-4-(trifluoromethyl)benzenamide
	methabenzthiazuron	1,3-dimethyl-3-(2-benzothiazolyl)urea
	metham	methylcarbamodithioic acid
20	methazole	2-(3,4-dichlorophenyl)-4-methyl-1,2,4-oxadiazolidine-3,5-dione
	methoxuron	N'-(3-chloro-4-methoxyphenyl)-N,N-dimethylurea
	metolachlor	2-chloro-N-(2-ethyl-6-methylphenyl)-N-(2-methoxy-1-methylethyl)acetamide
25	metribuzin	4-amino-6-(1,1-dimethylethyl)-3-(methylthio)-1,2,4-triazin-5(4H)-one
	metsulfuron methyl	2-[[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)amino]carbonyl]-amino]sulfonyl]benzoic acid, methyl ester
30	MH	1,2-dihydro-3,6-pyridazinedione
	molinate	S-ethyl hexahydro-1H-azepine-1-carbothioate
35		




	<u>Common Name</u>	<u>Chemical Name</u>
5	monolinuron	3-(p-chlorophenyl)-1-methoxy-1-methyl-urea
	monuron	N'-(4-chlorophenyl)-N,N-dimethylurea
	monuron TCA	Salt of monuron and TCA
10	MSMA	monosodium salt of MAA
	napropamide	N,N-diethyl-2-(1-naphthalenyloxy)-propanamide
	naptalam	2-[(1-naphthalenylamino)carbonyl]-benzoic acid
15	neburon	1-butyl-3-(3,4-dichlorophenyl)-1-methyl-urea
	nitralin	4-(methylsulfonyl)-2,6-dinitro-N,N-dipropylaniline
	nitrofen	2,4-dichloro-1-(4-nitrophenoxy)benzene
20	nitrofluorfen	2-chloro-1-(4-nitrophenoxy)-4-(trifluoromethyl)benzene
	norea	N,N-dimethyl-N'-(octahydro-4,7-methano-1H-inden-5-yl)urea 3 $\alpha$ ,-4 $\alpha$ ,5 $\alpha$ ,7 $\alpha$ ,7 $\alpha$ -isomer
25	norflurazon	4-chloro-5-(methylamino)-2-[3-(trifluoromethyl)phenyl]-3(2H)-pyridazinone
	oryzalin	4-(dipropylamino)-3,5-dinitro-benzenesulfonamide
30	oxadiazon	3-[2,4-dichloro-5-(1-methylethoxy)-phenyl]-5-(1,1-dimethylethyl)-1,3,4-oxadiazol-2(3H)-one
	oxyfluorfen	2-chloro-1-(3-ethoxy-4-nitrophenoxy)-4-(trifluoromethyl)benzene
35	paraquat	1,1'-dimethyl-4,4'-dipyridinium ion

	<u>Common Name</u>	<u>Chemical Name</u>
5	pebulate	S-propyl butylethylcarbamothioate
	pendimethalin	N-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine
	perfluidone	1,1,1-trifluoro-N-[2-methyl-4-(phenylsulfonyl)phenyl]methanesulfonamide
10	phenmedipham	3-[(methoxycarbonyl)amino]phenyl (3-methylphenyl)carbamate
	picloram	4-amino-3,5,6-trichloro-2-pyridine-carboxylic acid
15	PPG-1013	5-[2-chloro-4-(trifluoromethyl)-phenoxy]-2-nitroacetophenone oxime-O-acetic acid, methyl ester
	pretilachlor	$\alpha$ -chloro-2,6-diethyl-N-(2-propoxyethyl)acetanilide
20	procyzazine	2-[[4-chloro-6-(cyclopropylamino)-1,3,5-triazine-2-yl]amino]-2-methylpropane-nitrile
	profluralin	N-(cyclopropylmethyl)-2,6-dinitro-N-propyl-4-(trifluoromethyl)benzenamine
	prometon	6-methoxy-N,N'-bis(1-methylethyl)-1,3,5-triazine-2,4-diamine
25	prometryn	N,N'-bis(1-methylethyl)-6-(methylthio)-1,3,5-triazine-2,4-diamine
	pronamide	3,5-dichloro-N-(1,1-dimethyl-2-propynyl)benzamide
30	propachlor	2-chloro-N-(1-methylethyl)-N-phenylacetamide
	propanil	N-(3,4-dichlorophenyl)propanamide
	propazine	6-chloro-N,N'-bis(1-methylethyl)-1,3,5-triazine-2,4-diamine
35	propham	1-methylethyl phenylcarbamate

	<u>Common Name</u>	<u>Chemical Name</u>
5	prosulfalin	N-[[4-(dipropylamino)-3,5-dinitro-phenyl]sulfonyl]-S,S-dimethylsulfilimine
	prynachlor	2-chloro-N-(1-methyl-2-propynyl)acetanilide
10	pyrazolate	4-(2,4-dichlorobenzoyl)-1,3-dimethylpyrazol-5-yl-p-toluenesulphonate
	pyrazon	5-amino-4-chloro-2-phenyl-3(2H)-pyridazinone
15	pyrazosulfuron ethyl	ethyl 5-[3-(4,6-dimethoxypyrimidin-2-yl)ureadosulfonyl]-1-methylpyrazole-4-carboxylate
	quinclorac	3,7-dichloro-8-quinoline carboxylic acid
	quizalofop ethyl	(±)-2-[4-[(6-chloro-2-quinoxalinyloxy]phenoxy]propanoic acid, ethyl ester
20	secbumeton	N-ethyl-6-methoxy-N'-(1-methylpropyl)-1,3,5-triazine-2,4-diamine
	sethoxydim	2-[1-(ethoxyimino)butyl]-5-[2-(ethylthio)propyl]-3-hydroxy-2-cyclohexen-1-one
25	siduron	N-(2-methylcyclohexyl)-N'-phenylurea
	simazine	6-chloro-N,N'-diethyl-1,3,5-triazine-2,4-diamine
	SK-233	1-(α,α-dimethylbenzyl)-3-(4-methylphenyl)urea
30	sulfometuron methyl	2-[[[[(4,6-dimethyl-2-pyrimidinyl)-amino]carbonyl]amino]sulfonyl]benzoic acid, methyl ester
	TCA	trichloroacetic acid
35	tebuthiuron	N-[5-(1,1-dimethylethyl)-1,3,4-thiadiazol-2-yl]-N,N'-dimethylurea
	terbacil	5-chloro-3-(1,1-dimethylthyl)-6-methyl-2,4(1H,3H)-pyrimidinedione

	<u>Common Name</u>	<u>Chemical Name</u>
5	terbuchlor	N-(butoxymethyl)-2-chloro-N-[2-(1,1-dimethylethyl)-6-methylphenyl]-acetamide
	terbuthyl- azine	2-( <del>tert</del> -butylamino)-4-chloro-6-(ethylamino)- <del>s</del> -triazine
10	terbutol	2,6-di- <del>tert</del> -butyl-p-tolyl methylcarbamate
	terbutryn	N-(1,1-dimethylethyl)-N'-ethyl-6-(methylthio)-1,3,5-triazine-2,4-diamine
15	thiobencarb	S-[(4-chlorophenyl)methyl] diethylcarbamothioate
	triallate	S-(2,3,3-trichloro-2-propenyl) bis(1-methylethyl)carbamothioate
20	tribenuron methyl	2-[[[N-(4-methoxy-6-methyl-1,3,5-triazine-2-yl)-N-methylamino]carbonyl]amino]sulfonyl]benzoic acid, methyl ester
	triclopyr	[(3,5,6-trichloro-2-pyridinyl)-oxy]acetic acid
25	tridiphane	2-(3,5-dichlorophenyl)-2-(2,2,2-trichloroethyl)oxirane
	trifluralin	2,6-dinitro-N,N-dipropyl-4-(trifluoromethyl)benzenamine
	trimeturon	1-(p-chlorophenyl)-2,3,3-trimethylpseudourea
30	2,4-D	(2,4-dichlorophenoxy)acetic acid
	2,4-DB	4-(2,4-dichlorophenoxy)butanoic acid
	vernolate	S-propyl dipropylcarbamothioate
35	xylachlor	2-chloro-N-(2,3-dimethylphenyl)-N-(1-methylethyl)acetamide

Herbicidal properties of the subject compounds were discovered in a series of greenhouse tests. Test procedures and results follow.

COMPOUND TABLE

A = OR¹

<u>CMFD</u>	<u>R¹</u>	<u>R²</u>	<u>X</u>	<u>Y</u>	<u>m.p. (°C)</u>
1	CH₂CH₃	CH₂CN	CH₃	CH₃	169-175
2	CH₂CH₃	CH₂CN	CH₃	OCH₃	153-156
3	CH₂CH₃	CH₂CN	OCH₃	OCH₃	187-191
4	CH₂CH₃	CH₂CN	OCH₃	Cl	177-184
5	CH₃	CH₂CN	CH₃	CH₃	183-185
6	CH₃	CH₂CN	CH₃	OCH₃	168-170
7	CH₃	CH₂CN	OCH₃	OCH₃	186-194
8	CH₃	CH₂CN	OCH₃	Cl	186-194
9	CH₃	CH₂CN	OCH₃	OCF₂H	165-171
10	CH₃	CH₂CN	OCH₃	H	179-180(d)
11	CH₂CH₃	CH₂F	OCH₃	OCH₃	190-195
12	CH₃	CH₂F	OCH₃	OCH₃	168-170
13	CH₂CH₃	CH₂F	CH₃	OCH₃	177-179
14	CH₂CH₃	CH₂F	OCH₃	Cl	168-172
15	CH₃	CH₂F	OCH₃	Cl	165-172
16	CH₃	CH₂F	CH₃	OCH₃	162-163
17	CH₃	CH₂F	OCH₃	H	166-167(d)

<u>CMPD</u>	<u>R<sup>1</sup></u>	<u>R<sup>2</sup></u>	<u>X</u>	<u>Y</u>	<u>m.p. (°C)</u>
18	CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>2</sub> CN	CH <sub>3</sub>	OCH <sub>3</sub>	176-177
19	CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>2</sub> CN	OCH <sub>3</sub>	OCH <sub>3</sub>	196-199
20	CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>2</sub> CN	OCH <sub>3</sub>	Cl	196-200
21	CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>2</sub> CN	OCH <sub>3</sub>	H	174-175
22	CH <sub>3</sub>	CHF <sub>2</sub>	CH <sub>3</sub>	OCH <sub>3</sub>	165-167
23	CH <sub>3</sub>	CHF <sub>2</sub>	OCH <sub>3</sub>	OCH <sub>3</sub>	181-185
24	CH <sub>3</sub>	CHF <sub>2</sub>	OCH <sub>3</sub>	H	178-180(d)
25	CH <sub>3</sub>	CHF <sub>2</sub>	CH <sub>3</sub>	CH <sub>3</sub>	167-171
26	CH <sub>2</sub> CH <sub>3</sub>	CHF <sub>2</sub>	OCH <sub>3</sub>	H	148-155
27	CH <sub>2</sub> CH <sub>3</sub>	CHF <sub>2</sub>	OCH <sub>3</sub>	OCH <sub>3</sub>	192-194
28	CH <sub>2</sub> CH <sub>3</sub>	CHF <sub>2</sub>	CH <sub>3</sub>	OCH <sub>3</sub>	162-164
29	CH(CH <sub>3</sub> ) <sub>2</sub>	CHF <sub>2</sub>	CH <sub>3</sub>	OCH <sub>3</sub>	174-176
30	CH(CH <sub>3</sub> ) <sub>2</sub>	CHF <sub>2</sub>	OCH <sub>3</sub>	OCH <sub>3</sub>	177-180
31	CH(CH <sub>3</sub> ) <sub>2</sub>	CHF <sub>2</sub>	OCH <sub>3</sub>	Cl	186-196
32	CH <sub>2</sub> CH <sub>3</sub>	CHF <sub>2</sub>	CH <sub>3</sub>	CH <sub>3</sub>	180-192(d)
33	CH <sub>2</sub> CH <sub>3</sub>	CH <sub>2</sub> F	CH <sub>3</sub>	CH <sub>3</sub>	194-196
34	CH <sub>3</sub>	CH <sub>2</sub> F	CH <sub>3</sub>	CH <sub>3</sub>	180-182(d)
35	CH <sub>3</sub>	CHF <sub>2</sub>	CH <sub>3</sub>	CH <sub>3</sub>	167-169(d)
36	CH <sub>2</sub> CH <sub>3</sub>	CHF <sub>2</sub>	OCH <sub>3</sub>	Cl	173-177
37	CH <sub>3</sub>	CHFCH <sub>3</sub>	OCH <sub>3</sub>	OCH <sub>3</sub>	155-158
38	CH <sub>3</sub>	CHFCH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>	149-153(d)
39	CH <sub>3</sub>	CHFCH <sub>3</sub>	OCH <sub>3</sub>	Cl	134-137(d)
40	CH <sub>3</sub>	CHFCH <sub>3</sub>	CH <sub>3</sub>	OCH <sub>3</sub>	58-64(d)
41	CH <sub>2</sub> CH <sub>3</sub>	CHFCH <sub>3</sub>	OCH <sub>3</sub>	Cl	153-156(d)
43	CH <sub>3</sub>	CH <sub>2</sub> CN	CH <sub>3</sub>	CH <sub>2</sub> CH <sub>3</sub>	173-175
<u>CMPD</u>	<u>A</u>	<u>R<sup>2</sup></u>	<u>X</u>	<u>Y</u>	<u>m.p. (°C)</u>
42	N(CH <sub>3</sub> ) <sub>2</sub>	CHFCH <sub>3</sub>	OCH <sub>3</sub>	OCH <sub>3</sub>	156-158

**TEST A**

Seeds of barley (Hordeum vulgare),  
5 barnyardgrass (Echinochloa crus-galli), cheatgrass  
(Bromus secalinus), cocklebur (Xanthium  
pensylvanicum), corn (Zea mays), cotton (Gossypium  
hirsutum), crabgrass (Digitaria spp.), giant foxtail  
(Setaria faberi), morningglory (Ipomoea spp.), rice  
10 (Oryza sativa), sorghum (Sorghum bicolor), soybean  
(Glycine max), sugar beet (Beta vulgaris), velvetleaf  
(Abutilon theophrasti), wheat (Triticum aestivum),  
and wild oat (Avena fatua) and purple nutsedge  
(Cyperus rotundus) tubers were planted and treated  
15 preemergence with test chemicals dissolved in a  
non-phytotoxic solvent. At the same time, these crop  
and weed species were also treated with postemergence  
applications of test chemicals. Plants ranged in  
height from two to eighteen cm (two to three leaf  
20 stage) for postemergence treatments. Treated plants  
and controls were maintained in a greenhouse for  
approximately sixteen days, after which all species  
were compared to controls and visually evaluated.  
Plant response ratings, summarized in Table A, are  
25 based on a scale of 0 to 10 where 0 is no effect and  
10 is complete control. A dash (-) response means no  
test result.

30

35

Tabl A

Rat (50 g/ha)	COMPOUND																																			
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	36				
POSTEMERGENCE																																				
Barley	9	9	8	3	9	9	9	8	7	0	2	5	9	0	6	9	2	7	9	8	0	9	3	0	9	0	0	8	9	8	6	0				
Barnyardgrass	9	9	9	9	10	9	9	9	8	9	9	10	9	9	10	7	9	9	8	1	9	9	7	9	7	8	9	9	9	7	7					
Cheatgrass	9	10	10	9	9	9	9	9	3	9	9	9	9	10	9	9	8	9	9	8	0	9	9	9	9	8	7	9	9	6	6	9				
Cockl bur	10	10	10	10	9	10	10	10	9	9	9	10	10	9	10	10	9	10	10	10	9	10	10	9	10	9	10	10	10	10	10	9				
Corn	9	9	9	6	5	9	9	3	6	0	9	9	9	7	7	10	4	9	9	8	2	10	9	0	9	0	5	10	9	8	3	2				
Cotton	10	9	9	9	10	9	9	9	9	10	9	10	9	9	10	10	9	9	9	9	10	9	9	9	9	9	9	10	9	9	9	10				
Crabgrass	-	9	8	5	9	10	6	8	7	2	9	8	6	8	6	9	0	6	3	1	0	3	1	4	9	0	0	9	4	5	4	0				
Giant f xtail	9	9	9	8	9	10	9	9	6	5	5	9	8	3	8	9	3	9	9	3	0	10	6	4	9	0	2	8	7	8	6	1				
M rningglory	9	10	10	8	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	9	10	10	10	9	10	10	10	10	10	10	9				
Nutsedge	9	10	10	10	-	10	10	10	10	6	10	-	10	10	10	10	9	9	10	-	-	10	10	9	10	9	8	9	9	10	10	0				
Ric	9	9	9	9	9	9	9	9	4	9	9	8	7	9	9	4	9	9	9	5	9	9	8	9	4	3	8	9	8	7	5					
S rghum	9	9	9	9	9	9	10	9	9	9	9	9	9	8	9	9	9	9	9	8	9	-	9	-	6	4	9	9	6	7	6					
S yb an	9	9	9	7	9	9	9	9	9	9	9	9	9	6	9	9	9	9	9	7	9	9	9	9	9	9	9	9	9	7	9					
Sugar b et	9	9	10	10	9	10	9	9	10	10	10	9	10	9	9	9	9	10	10	9	9	10	10	9	10	10	10	10	10	9	9	10				
V lvetleaf	10	10	10	9	10	10	9	10	10	9	9	10	10	9	9	10	9	9	9	-	-	9	9	10	9	9	-	9	9	10	9					
Wheat	9	9	9	7	9	9	9	7	2	7	5	9	2	8	9	0	9	9	3	0	9	3	0	9	0	0	8	6	2	0	0					
Wild at	9	9	7	5	9	9	7	8	6	4	3	3	8	4	6	9	2	8	5	3	0	7	4	2	9	0	0	6	5	0	3	3				



Table A

	COMPOUND																																			
Rate (50 g/ha)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	36				
PREEMERGENCE																																				
Barley	9	8	9	6	9	9	9	9	3	0	3	0	8	2	4	9	0	8	6	2	0	7	3	2	9	1	3	7	7	0	1	2				
Barnyardgrass	9	9	9	8	9	9	9	9	6	0	5	9	9	0	9	9	4	9	9	9	1	9	7	2	9	0	2	8	8	7	4	5				
Ch atgrass	9	9	9	7	9	9	9	7	4	2	8	8	8	6	8	9	4	8	8	8	6	8	7	5	8	5	6	7	9	7	7	5				
Cocklebur	9	9	9	8	9	8	9	5	7	1	9	-	9	7	7	9	4	9	9	6	2	9	8	6	9	6	9	8	8	8	6	8				
Corn	9	9	9	7	8	9	8	1	0	0	0	3	9	0	2	9	0	9	8	8	0	8	2	0	9	3	5	9	7	1	0	3				
Cott n	9	9	9	3	9	9	9	7	0	2	3	2	8	1	7	9	6	7	8	7	0	4	8	4	9	8	1	7	6	4	-	5				
Crabgrass	8	9	8	6	10	9	9	9	7	0	4	4	9	2	9	9	2	6	4	2	0	7	4	6	10	0	5	7	6	3	0	2				
Giant foxtail	9	9	8	6	9	9	9	8	3	2	5	5	7	0	6	9	3	8	6	7	0	9	5	2	9	0	2	6	7	4	2	2				
Morningglory	9	8	9	9	9	9	9	9	2	0	8	2	8	9	9	7	9	9	6	2	9	9	4	9	3	9	7	9	6	9	9	9				
Nutsedge	10	8	8	8	10	10	9	9	0	0	10	9	0	8	9	0	8	10	10	0	8	0	4	9	10	0	-	6	10	10	0	0				
Ric	9	9	9	9	10	9	9	9	8	0	5	8	9	7	8	9	7	10	9	9	5	9	8	8	10	6	5	9	8	5	4	4				
Sorghum	10	9	9	9	10	9	9	9	9	6	3	4	9	7	9	9	8	9	9	9	7	9	7	8	10	4	2	9	9	2	4	6				
Soybean	9	9	9	5	9	9	9	8	4	3	1	4	9	0	1	8	4	9	9	3	1	7	7	7	8	6	7	8	8	3	5	5				
Sugar beet	9	9	9	9	9	9	9	9	9	8	6	8	9	8	9	9	7	9	9	9	9	9	9	8	9	9	9	9	9	8	9	8				
Velvetleaf	9	9	8	8	9	9	9	8	9	3	7	7	9	6	9	9	9	9	5	6	5	8	7	8	9	7	7	9	9	8	5	9				
Wheat	9	8	9	7	9	9	9	8	0	0	0	9	0	9	0	2	9	0	9	8	8	0	8	0	9	0	0	9	6	0	0	0				
Wild at	8	8	5	5	8	8	7	6	0	2	0	0	8	0	3	8	0	8	3	3	2	6	2	0	8	0	0	4	5	0	0	0				

Tabl A

Rate (10 g/ha)	COMPOUND																																			
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	36				
POSTEMERGENCE																																				
Barley	9	8	5	2	9	9	9	3	2	0	0	0	8	0	0	9	0	8	5	2	0	9	2	0	9	0	0	7	7	6	2	0				
Barnyardgrass	9	9	9	8	9	9	9	9	8	6	9	9	9	7	9	10	6	9	9	3	0	9	7	6	8	2	8	9	8	9	4	5				
Cheatgrass	9	9	9	6	9	9	9	9	2	3	9	9	9	8	9	9	6	8	6	5	0	9	9	7	10	8	8	4	6	6	4	5				
Cocklebur	9	9	10	9	9	9	10	9	9	10	10	9	10	9	10	9	9	9	10	9	10	10	9	10	10	10	10	10	10	9	10	10				
C rn	7	9	9	3	0	9	9	0	2	0	9	9	9	2	3	9	2	9	9	6	0	9	8	0	9	0	2	9	9	5	0	0				
Cotton	9	9	9	9	9	9	9	9	9	9	9	9	9	9	10	10	9	9	9	9	9	10	9	9	9	9	10	10	9	9	9	9				
Crabgrass	0	2	2	0	6	5	2	2	0	0	4	9	8	2	2	4	0	0	1	0	0	2	0	0	9	0	0	9	2	2	0	0				
Giant f xtall	8	7	5	2	9	9	9	4	2	0	3	7	6	0	6	9	2	6	4	0	0	8	3	0	9	0	1	5	5	7	2	0				
Morningglory	9	9	10	10	6	10	10	10	9	9	10	10	9	10	10	10	9	10	9	9	8	10	10	10	10	10	10	10	10	9	9	10				
Nutsedge	9	9	-	9	9	-	-	-	-	0	9	10	10	10	10	10	9	9	8	9	0	10	10	9	10	10	9	8	10	9	0					
Rice	9	9	8	8	9	9	9	9	7	2	3	9	9	1	7	9	3	9	9	9	2	8	9	7	9	0	2	6	5	4	4	0				
S rghum	9	9	9	8	9	9	9	9	9	9	9	9	9	5	9	9	8	9	9	8	5	9	7	6	9	0	5	7	6	6	4	4				
S yb an	9	9	9	5	9	9	9	8	8	-	9	9	9	5	5	9	8	9	9	3	2	8	9	9	10	9	9	8	9	4	8					
Sugar be t	9	9	9	9	9	8	9	9	9	9	10	9	10	10	9	9	9	9	9	8	10	10	10	10	10	10	10	9	9	8	10					
Velv tleaf	9	10	9	8	9	9	10	9	9	8	9	10	9	9	9	9	8	9	9	-	8	-	9	9	10	9	9	7	9	9	9					
Wheat	9	9	6	2	9	9	9	2	0	0	0	0	3	0	2	9	0	9	8	0	0	5	0	0	9	0	0	3	0	0	0	0				
Wild at	9	8	4	2	9	8	5	3	3	2	0	2	2	0	4	8	0	7	0	0	0	4	2	0	9	0	0	2	0	0	3	0				

Table A

Rate (10 g/ha)	COMPOUND																																			
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	36				
PREMERGENCE																																				
Barl y	8	8	7	2	9	9	8	7	0	0	0	0	6	0	0	8	0	6	0	0	0	5	3	0	8	0	0	3	2	0	0	0	0	0	0	
Barnyardgrass	7	9	7	7	9	9	9	8	0	0	0	4	4	0	7	9	3	9	5	3	0	7	5	0	8	0	0	0	3	2	0	0	0	0	0	
Cheatgrass	8	8	8	7	9	9	8	7	2	0	2	5	7	0	6	8	2	7	7	5	0	5	2	2	8	2	0	3	7	2	2	0	0	0	0	
C cklebur	7	7	7	5	9	8	8	5	0	0	-	3	9	0	7	4	3	8	5	7	2	7	5	-	8	2	3	2	6	3	-	1	0	0	0	
Corn	9	9	8	3	6	6	8	0	0	0	0	2	7	0	0	7	2	8	5	6	0	3	0	0	9	0	0	0	2	0	0	0	0	0	0	
Cotton	7	4	2	3	8	8	9	2	0	0	0	0	3	0	2	8	4	7	2	3	0	2	0	0	8	1	0	1	5	2	0	0	0	0	0	
Crabgrass	7	7	4	2	9	9	7	9	0	0	0	4	2	0	7	9	0	2	2	0	0	5	4	2	8	0	0	2	0	0	0	0	0	0	0	
Giant foxtail	9	8	7	1	9	9	9	2	3	0	0	2	3	0	2	9	3	7	2	3	0	6	0	0	7	0	0	2	2	2	0	0	0	0	0	
Morninggl ry	3	4	9	8	7	9	9	8	2	0	0	4	3	0	4	7	6	7	0	0	0	9	7	2	8	0	3	2	2	1	1	9	0	0	0	
Nutsedg	9	8	5	3	10	9	10	0	7	0	0	0	2	0	10	7	-	7	5	0	0	0	0	0	0	0	0	0	0	10	6	4	0	0		
Rice	-	9	9	9	9	9	8	9	5	0	0	7	8	0	7	9	3	9	8	7	0	6	0	3	9	2	0	4	4	0	0	0	0	0	0	
Sorghum	9	10	9	9	10	9	9	9	3	2	0	0	8	0	7	9	3	9	9	8	2	7	6	3	9	3	0	3	5	2	2	0	0	0	0	
Soybean	8	9	6	5	9	8	6	3	1	0	1	3	7	0	0	7	2	7	1	1	0	4	3	5	9	2	2	3	4	2	0	0	0	0	0	
Sugar beet	9	9	9	9	9	9	9	7	8	4	2	6	7	0	8	9	4	8	5	5	2	9	6	3	8	9	4	9	8	4	7	8	0	0	0	
velvetl af	9	7	4	5	9	8	8	7	2	2	0	5	9	0	7	9	6	8	2	7	2	3	3	4	3	1	5	2	8	3	2	0	0	0	0	
Wheat	9	9	8	5	8	9	8	7	0	0	0	0	5	0	0	9	0	8	4	2	0	3	0	0	6	0	0	0	0	0	0	0	0	0	0	
Wild at	8	6	4	0	8	8	3	0	0	0	0	0	5	0	0	6	0	7	0	0	2	0	0	3	0	0	0	2	0	0	0	0	0	0	0	

**TEST B**

- Seeds of barley (Hordeum vulgare),  
5 barnyardgrass (Echinochloa crus-galli), blackgrass  
(Alopecurus myosuroides), cheatgrass (Bromus  
secalinus), chickweed (Stellaria media), cocklebur  
(Xanthium pensylvanicum), corn (Zea mays), cotton  
(Gossypium hirsutum), crabgrass (Digitaria spp.),  
10 bedstraw (Galium aparine), giant foxtail (Setaria  
faberii), lambsquarters (Chenopodium album),  
morningglory (Ipomoea hederacea), rape (Brassica  
napus), rice (Oryza sativa), sorghum (Sorghum  
bicolor), soybean (Glycine max), sugar beet (Beta  
15 vulgaris), velvetleaf (Abutilon theophrasti), wheat  
(Triticum aestivum), wild buckwheat (Polygonum  
convolvulus), and wild oat (Avena fatua) and purple  
nutsedge (Cyperus rotundus) tubers were planted and  
treated preemergence with test chemicals dissolved in  
20 a non-phytotoxic solvent. At the same time, these  
crop and weed species were also treated with  
postemergence applications of test chemicals. Plants  
ranged in height from two to eighteen cm (one to four  
leaf stage) for postemergence treatments. Treated  
25 plants and controls were maintained in a greenhouse  
for twelve to sixteen days, after which all species  
were compared to controls and visually evaluated.  
Plant response ratings, summarized in Table B, are  
based on a scale of 0 to 10 where 0 is no effect and  
30 10 is complete control. A dash (-) response means no  
test result.

59

Table B

	COMPOUND
Rate (200 g/ha)	43
POSTEMERGENCE	
Barley	8
Barnyardgrass	10
Bedstraw	9
Blackgrass	10
Cheatgrass	9
Chickweed	10
Corn	9
Cotton	9
Crabgrass	6
Giant foxtail	9
Lambsquarters	9
Morningglory	10
Nutsedge	10
Rape	10
Rice	9
Sorghum	9
Soybean	9
Sugar beet	9
Velvetleaf	9
Wheat	9
Wild buckwheat	9
Wild oat	8

60

## Tabl B

	COMPOUND
Rate (200 g/ha)	43
PREEMERGENCE	
Barley	8
Barnyardgrass	9
Bedstraw	9
Blackgrass	9
Cheatgrass	8
Chickweed	9
Corn	9
Cotton	8
Crabgrass	9
Giant foxtail	9
Lambsquarters	9
Morningglory	9
Nutsedge	10
Rape	9
Rice	10
Sorghum	9
Soybean	9
Sugar beet	9
Velvetleaf	9
Wheat	9
Wild buckwheat	9
Wild oat	8

Tabl B

	COMPOUND											
Rate (50 g/ha)	32	33	34	35	37	38	39	40	41	42	43	
POSTEMERGENCE												
Barley	9	9	10	10	7	9	5	9	5	9	7	
Barnyardgrass	9	10	10	9	9	9	10	9	9	10	9	
Bedstraw	10	9	9	10	9	10	10	10	10	10	7	
Blackgrass	9	10	10	9	9	9	9	9	8	9	9	
Cheatgrass	9	9	10	9	9	9	9	9	9	9	8	
Chickweed	10	-	-	10	10	10	9	10	9	10	9	
Cocklebur	10	10	10	10	10	10	10	10	-	10	-	
Corn	10	10	9	10	10	10	9	9	9	10	7	
Cotton	9	9	10	10	9	10	10	9	10	9	9	
Crabgrass	9	-	-	9	4	10	5	4	0	5	4	
Giant foxtail	9	9	10	9	9	10	9	9	6	9	9	
Lambsquarters	10	10	10	10	10	10	9	9	10	10	9	
Morningglory	9	9	9	9	10	10	9	10	10	10	10	
Nutsedge	10	10	10	9	10	10	10	10	9	-	10	
Rape	10	9	9	9	9	9	9	10	10	9	9	
Rice	9	9	10	9	9	10	9	9	8	9	9	
Sorghum	9	9	10	9	9	10	9	9	8	10	9	
Soybean	9	9	9	9	9	9	9	9	9	9	9	
Sugar beet	10	9	9	10	9	9	9	10	10	10	9	
Velvetleaf	9	10	10	10	10	10	10	9	9	10	9	
Wheat	9	9	9	9	8	9	4	9	5	9	8	
Wild buckwheat	10	10	10	10	10	10	10	10	10	9	7	
Wild oat	9	9	10	9	7	9	8	9	7	6	5	

Table B

	COMPOUND											
Rate (50 g/ha)	32	33	34	35	37	38	39	40	41	42	43	
PREEMERGENCE												
Barley	9	9	9	9	2	9	1	2	2	9	5	
Barnyardgrass	9	9	9	9	7	9	5	7	5	9	9	
Bedstraw	7	8	8	7	7	9	9	8	8	9	8	
Blackgrass	9	9	9	8	5	9	4	9	8	9	8	
Cheatgrass	9	9	10	9	8	9	8	9	8	9	8	
Chickweed	10	-	-	10	9	10	10	-	9	10	8	
Cocklebur	9	9	9	-	6	8	9	-	-	9	-	
Corn	9	9	9	9	7	9	3	8	6	9	7	
Cotton	9	9	9	9	3	9	8	8	0	8	5	
Crabgrass	7	9	9	8	5	9	2	6	2	7	7	
Giant foxtail	7	6	8	9	5	8	6	7	2	9	8	
Lambsquarters	10	10	-	10	9	9	10	10	9	10	9	
Morningglory	8	6	9	10	9	9	9	9	9	9	9	
Nutsedge	9	10	9	10	7	10	9	0	9	10	10	
Rape	10	9	9	9	9	10	9	9	9	9	8	
Rice	10	9	10	9	8	10	8	7	8	9	9	
Sorghum	9	9	9	9	9	10	8	7	0	9	9	
Soybean	9	9	9	9	9	9	6	8	2	9	7	
Sugar beet	9	8	8	9	9	9	9	9	9	9	9	
Velvetleaf	9	9	9	9	8	9	7	9	7	9	8	
Wheat	9	9	9	9	0	9	2	7	2	9	9	
Wild buckwheat	8	8	8	8	5	6	1	5	5	9	8	
Wild oat	9	9	9	9	1	9	1	2	7	2	7	



Table B

	COMPOUND										
Rate (10 g/ha)	32	33	34	35	37	38	39	40	41	42	
POSTEMERGENCE											
Barley	9	9	9	9	4	9	4	9	0	9	
Barnyardgrass	8	10	10	9	9	9	9	9	8	10	
Bedstraw	6	7	7	8	10	9	10	10	9	9	
Blackgrass	9	9	9	8	8	9	7	9	9	10	
Cheatgrass	9	9	9	9	9	9	7	9	5	9	
Chickweed	10	9	-	10	10	9	10	10	8	9	
Cocklebur	10	9	10	10	10	10	10	9	-	-	
Corn	9	9	9	9	9	10	1	9	4	9	
Cotton	9	9	9	9	9	9	9	9	9	9	
Crabgrass	-	-	-	7	-	9	2	3	0	0	
Giant foxtail	8	7	10	9	5	9	5	8	3	8	
Lambsquarters	9	9	10	10	10	9	10	9	9	9	
Morningglory	6	3	8	9	10	10	9	9	10	10	
Nutsedge	9	10	10	9	10	10	10	9	9	8	
Rape	10	9	9	9	9	9	9	10	10	9	
Rice	9	9	10	9	7	9	3	8	2	9	
Sorghum	9	9	10	9	9	10	8	9	2	9	
Soybean	9	9	9	9	9	9	9	9	8	9	
Sugar beet	10	5	9	9	10	9	9	9	10	9	
Velvetleaf	9	9	10	9	10	9	9	9	9	9	
Wheat	9	9	9	9	3	9	2	7	1	9	
Wild buckwheat	-	7	10	9	9	9	10	9	9	7	
Wild oat	9	9	10	9	3	9	2	5	2	2	

Tabl B

	COMPOUND										
Rate (10 g/ha)	32	33	34	35	37	38	39	40	41	42	
PREEMERGENCE											
Barley	7	8	8	2	0	3	0	1	0	3	
Barnyardgrass	3	4	9	7	3	7	0	3	2	7	
Bedstraw	7	5	8	0	3	5	0	3	1	3	
Blackgrass	8	8	9	7	2	7	3	6	0	7	
Cheatgrass	9	9	9	9	7	9	7	8	7	8	
Chickweed	10	-	-	-	10	9	10	-	8	10	
Cocklebur	7	4	7	-	0	8	7	0	-	6	
Corn	9	9	9	8	1	7	0	2	0	0	
Cotton	5	9	8	9	0	6	8	6	0	3	
Crabgrass	2	6	6	3	1	9	0	2	0	0	
Giant foxtail	3	6	2	5	0	7	0	5	0	4	
Lambsquarters	9	-	9	9	10	9	10	10	6	9	
Morningglory	4	4	5	9	3	9	6	3	0	6	
Nutsedge	5	9	9	9	0	10	7	3	0	0	
Rape	9	9	9	9	8	9	2	2	3	9	
Rice	9	9	9	9	4	9	6	2	3	6	
Sorghum	1	9	8	9	0	8	0	1	0	3	
Soybean	7	6	6	7	3	6	2	2	0	6	
Sugar beet	8	7	7	8	8	8	9	7	8	8	
Velvetleaf	7	2	9	7	7	8	5	5	2	3	
Wheat	7	8	9	7	0	6	0	2	0	6	
Wild buckwheat	5	6	8	3	2	3	0	0	2	2	
Wild oat	5	8	9	6	0	4	0	3	4	0	

**TEST C**

Seeds of barley (Hordeum vulgare),  
5 barnyardgrass (Echinochloa crus-galli), blackgrass  
(Alopecurus myosuroides), chickweed (Stellaria  
media), cocklebur (Xanthium pensylvanicum), corn (Zea  
mays), cotton (Gossypium hirsutum), crabgrass  
(Digitaria spp.), downy brome (Bromus tectorum),  
10 giant foxtail (Setaria faberi), green foxtail  
(Setaria viridis), jimsonweed (Datura stramonium),  
johnsongrass (Sorghum halepense), lambsquarters  
(Chenopodium album), morningglory (Ipomoea spp.),  
rape (Brassica napus), rice (Oryza sativa), sicklepod  
15 (Cassia obtusifolia), soybean (Glycine max), sugar  
beet (Beta vulgaris), teaweed (Sida spinosa),  
velvetleaf (Abutilon theophrasti), wheat (Triticum  
aestivum), wild buckwheat (Polygonum convolvulus),  
and wild oat (Avena fatua) and purple nutsedge  
20 (Cyperus rotundus) tubers were planted and treated  
preemergence with test chemicals dissolved in a  
non-phytotoxic solvent. At the same time, these crop  
and weed species were also treated with postemergence  
applications of test chemicals. Plants ranged in  
25 height from two to eighteen cm (two to three leaf  
stage) for postemergence treatments. Treated plants  
and controls were maintained in a greenhouse for  
approximately 24 days, after which all species were  
compared to controls and visually evaluated. Plant  
30 response ratings, summarized in Table C, are reported  
on a 0 to 10 scale where 0 is no effect and 10 is  
complete control. A dash (-) response means no test  
result.

Table C

Rate (250 g/ha)	COMPOUND	
	4	10
POSTEMERGENCE		
Barley	7	-
Blackgrass	9	5
Chickweed	10	10
Cocklebur	10	10
Corn	8	2
Cotton	10	9
Crabgrass	7	3
Downy brome	7	5
Giant foxtail	7	7
Green foxtail	8	7
Jimsonweed	10	10
Johnsongrass	9	10
Lambsquarters	10	10
Morningglory	10	10
Nutsedge	10	4
Rape	10	10
Rice Dry Seed	10	9
Sicklepod	10	8
Soybean	9	10
Sugar beet	10	10
Tea weed	9	7
Velvetleaf	10	10
Wheat	8	3
Wild buckwheat	9	10
Wild oat	9	2
Barnyardgrass	10	9

Table C

Rate (250 g/ha)	COMPOUND	
	4	10
PREEMERGENCE		
Barley	7	0
Blackgrass	9	-
Chickweed	7	7
Cocklebur	10	6
Corn	9	0
Cotton	7	4
Crabgrass	9	10
Downy brome	9	6
Giant foxtail	8	7
Green foxtail	9	7
Jimsonweed	9	8
Johnsongrass	9	8
Lambsquarters	10	-
Morningglory	9	8
Nutsedge	9	0
Rape	10	9
Rice Dry Seed	10	9
Sicklepod	9	5
Soybean	8	3
Sugar beet	9	9
Teaweed	8	6
Velvetleaf	9	10
Wheat	9	0
Wild buckwheat	9	7
Wild oat	8	-
Barnyardgrass	10	6

Tabl C

		COMPOUND																																	
Rate (62 g/ha)		1	2	3	4	5	6	7	8	9	10	11	12	13	15	16	17	18	19	20	22	23	24	25	26	27	28	29	30	31	36				
POSTEMERGENCE																																			
Barley		9	9	8	6	10	10	9	8	-	-	3	4	7	4	9	2	6	4	0	10	4	5	10	5	4	9	8	2	3	5				
Blackgrass		9	10	9	9	10	10	10	9	10	5	10	7	9	6	8	4	9	8	7	10	6	5	10	6	7	10	10	9	5	3				
Chickweed		10	10	10	9	10	10	10	10	10	9	10	9	10	8	10	10	8	10	0	10	10	10	10	6	10	10	10	10	6	9				
Cockl bur		10	10	10	9	10	10	10	9	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10				
Corn		9	10	10	7	9	10	10	2	6	0	10	10	10	6	10	0	10	10	7	10	10	4	10	0	7	10	10	10	6	7				
C tt n		10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	9	10	10	10	10	10	10	10	10	10	10	10	10	10				
Crabgrass		8	5	8	5	9	7	7	8	5	2	7	9	6	3	9	4	6	4	3	10	-	6	10	0	4	7	5	0	0	6				
D wny brome		9	9	8	7	10	10	9	7	5	4	3	2	7	6	9	3	6	5	0	10	4	0	10	0	0	8	9	0	3	0				
Giant foxtail		10	9	10	7	10	10	10	8	5	5	4	7	7	7	10	2	8	6	3	10	6	5	10	0	0	5	0	0	4	2				
Gr en foxtail		9	8	8	7	10	10	10	8	2	6	5	7	7	6	10	4	7	6	3	10	7	7	10	0	2	8	4	0	4	2				
Jims nwe d		9	10	9	9	10	10	10	9	9	10	10	10	10	10	10	10	10	9	10	10	10	10	10	10	10	10	10	10	10	10				
Johnsongrass		10	10	10	7	10	10	10	10	8	10	7	9	10	9	10	6	7	9	7	10	6	8	10	7	2	10	5	4	7	5				
Lambsquarters		10	10	10	8	10	10	10	10	10	10	10	10	10	10	10	8	-	-	-	10	10	10	-	-	-	-	-	-	-	10				
Morningglory		10	10	10	9	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10				
Nutsedge		10	10	10	10	10	10	10	10	10	4	10	10	10	10	10	10	10	10	10	10	10	10	10	5	10	10	10	6	10	10				
Rap		10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10				
Rice Dry Seed		10	10	10	9	10	10	10	9	6	5	7	6	7	9	6	8	7	6	8	7	7	10	9	10	6	4	8	9	5	5				
Sicklepod		10	10	10	8	10	10	10	9	10	6	10	10	9	10	10	9	10	10	8	10	10	10	10	6	10	10	10	10	7	10				
Soybean		10	10	10	9	9	10	10	10	10	10	10	10	10	9	9	10	10	9	10	10	9	10	10	10	10	10	10	10	10	10				
Sugar beet		10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	8	10	10	10	10	10	10	10	10	10	10				
T awe d		9	9	8	8	10	9	8	9	6	5	7	9	9	9	10	8	9	7	8	10	10	10	10	9	6	10	7	8	8	10				
Velvetl af		10	10	10	9	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	9	10	9				
Wheat		9	8	9	5	10	10	9	9	6	0	1	2	8	4	9	2	8	6	3	10	5	0	10	0	3	7	6	0	0	0				
Wild buckwheat		10	10	10	9	10	10	10	10	10	9	10	10	10	10	9	9	10	10	8	10	10	10	10	10	9	10	10	10	9	9				
Wild at		10	9	9	5	10	10	9	9	0	0	1	0	6	8	8	0	6	5	3	8	0	0	10	3	4	5	6	0	0	4				
Barnyardgrass		10	10	10	9	10	10	10	10	10	8	8	10	10	10	10	7	8	8	7	10	10	10	10	0	6	10	9	8	4	8				

Table C

COMPOUND																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																										
Rat (62 g/ha)	1	2	3	4	5	6	7	8	9	10	11	12	13	15	16	17	18	19	20	22	23	24	25	26	27	28	29	30	31	36																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																												

Tabl C

	COMPOUND																																	
	1	2	3	4	5	6	7	8	9	10	11	12	13	15	16	17	18	19	20	22	23	24	25	26	27	28	29	30	31	36				
Rat (16 g/ha) POSTEMERGENCE																																		
Barley	8	8	7	4	9	9	7	4	-	-	0	3	6	2	8	0	4	0	0	9	3	3	10	3	0	6	5	0	0	0				
Blackgrass	9	9	9	8	9	10	10	8	9	0	5	5	8	4	7	2	8	6	5	9	5	4	8	4	3	10	9	5	2	3				
Chickwe d	10	10	10	7	9	10	10	8	9	8	8	9	9	8	10	7	7	7	0	10	10	7	10	5	9	9	10	10	0	9				
Cocklebur	9	9	10	8	10	9	10	9	10	9	10	10	10	10	10	10	10	10	9	10	10	10	10	10	10	10	10	10	-	10				
C rn	8	9	9	4	3	9	9	0	4	0	10	10	9	2	10	0	8	10	5	10	9	2	10	0	4	10	9	7	1	4				
C tton	9	9	10	9	10	9	10	9	10	10	8	10	10	9	9	7	9	10	8	10	10	9	10	6	10	10	9	7	6	10				
Crabgrass	7	5	5	3	7	7	3	5	4	0	6	6	5	3	7	2	4	2	3	10	10	6	10	-	3	6	3	0	0	6				
Downy brome	7	8	7	5	10	9	9	4	4	3	0	0	6	4	8	2	4	3	0	9	4	0	10	0	0	7	3	0	0	0				
Giant f xtall	9	9	7	4	10	10	9	6	4	4	3	6	5	4	9	0	6	3	0	8	5	4	10	0	0	4	-	0	1	0				
Green foxtail	8	7	5	3	10	9	9	7	0	4	4	6	5	4	9	2	6	3	0	10	6	4	10	0	0	5	0	0	0	0				
Jims nw ed	8	10	9	7	7	10	9	9	8	6	7	8	10	10	10	6	7	8	6	10	10	7	10	6	10	10	8	10	9	10				
J hnsongrass	10	9	9	7	10	10	10	8	8	6	6	8	8	7	9	5	6	6	3	10	6	5	10	2	2	8	4	3	6	2				
Lambsquarters	9	10	8	8	10	10	10	10	10	10	10	10	10	5	-	5	-	-	-	10	-	10	10	-	-	-	-	-	-	8				
M rningglory	9	10	10	9	9	10	10	10	10	10	10	10	10	10	10	10	10	8	9	10	10	10	9	10	10	10	10	10	10	10				
Nutsedge	10	10	10	10	10	10	10	10	10	3	10	10	10	9	10	10	7	10	10	-	10	10	10	4	10	10	8	-	10	9				
Rap	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10				
Rice Dry S ed	10	10	9	8	10	10	10	9	8	5	3	6	5	6	7	4	7	6	6	9	9	8	10	4	2	7	6	3	0	3				
Sickl p d	10	10	10	8	9	10	10	9	9	4	10	10	9	10	10	4	9	9	7	10	10	9	10	0	10	10	8	10	2	-				
S ybean	9	10	10	8	9	9	10	8	10	10	10	10	10	7	9	9	10	10	8	10	10	10	10	10	10	10	10	10	9	3				
Sugar beet	10	10	10	9	10	10	10	10	10	10	10	10	10	10	10	10	10	10	6	10	10	10	10	10	10	10	10	10	10	9				
Teaweed	8	7	5	7	8	9	5	8	6	3	5	7	8	9	10	6	7	5	7	10	9	6	10	8	6	8	6	6	4	7				
V lvetleaf	10	10	10	9	10	10	10	10	10	10	10	10	10	10	10	10	7	10	9	8	10	10	7	10	9	7	10	10	7	9				
Wheat	8	6	8	3	9	10	9	8	0	0	0	0	7	1	8	0	4	3	0	10	4	0	10	0	0	5	4	0	0	0				
Wild buckwheat	10	10	10	9	10	9	10	10	8	7	9	10	9	8	9	5	9	7	6	10	9	9	10	10	8	10	9	6	7	7				
Wild oat	10	9	8	4	10	10	6	9	0	0	0	0	5	4	5	0	4	2	0	7	0	0	9	0	0	4	3	0	0	2				
Barnyardgrass	10	10	9	9	10	10	10	9	8	6	7	10	9	10	10	4	8	6	3	10	10	9	10	0	6	8	8	8	4	3				



Table C

Rate (16 g/ha) PREEMERGENCE		COMPOUND																																			
		1	2	3	4	5	6	7	8	9	10	11	12	13	15	16	17	18	19	20	22	23	24	25	26	27	28	29	30	31	36						
Barley	8	8	3	2	10	9	2	2	0	0	0	2	0	7	2	6	0	6	5	3	5	3	3	0	5	0	0	6	6	3	3	2					
Blackgrass	8	8	7	5	10	10	8	8	-	-	-	3	3	7	3	7	3	9	8	7	6	0	0	5	5	4	7	6	3	3	3						
Chickweed	6	7	7	5	8	9	9	4	8	-	8	8	8	6	9	4	8	7	3	8	8	8	6	6	2	8	6	8	7	3	5						
C ckl bur	8	8	-	8	9	8	4	5	0	0	0	7	8	7	-	7	3	10	9	8	8	7	5	8	7	-	10	8	8	7	8						
C rn	8	5	3	2	2	7	4	1	0	0	0	3	7	0	4	0	8	0	5	6	4	0	10	0	7	8	5	0	0	4							
Cotton	9	5	5	3	6	7	2	8	3	0	6	8	8	4	7	3	4	4	3	8	3	6	8	6	5	9	5	3	3	7							
Crabgrass	9	9	8	8	8	9	7	8	8	10	7	8	5	7	9	3	5	6	3	6	5	0	7	5	8	8	7	6	7	5							
Downy brome	8	8	7	3	9	10	8	4	3	0	0	3	8	3	7	0	8	6	0	4	3	0	7	3	2	6	6	3	3	5							
Giant f xtail	8	9	7	4	9	9	8	7	3	0	3	3	5	0	6	0	8	6	3	9	0	0	10	2	2	7	4	0	0	3							
Gr en f xtail	9	9	7	7	9	9	8	7	3	0	3	4	7	3	7	3	9	6	4	8	6	5	10	3	0	6	6	3	3	4							
Jimsonweed	8	8	5	3	8	9	9	4	6	0	8	8	8	8	8	3	8	7	6	9	8	7	9	5	8	9	8	6	8	8							
Johnsongrass	9	8	7	4	9	8	8	8	6	3	5	5	8	6	8	3	9	8	8	9	3	5	10	6	0	8	6	3	5	2							
Lambsquarters	10	9	9	9	10	10	10	10	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	7	-	-	-	-	-							
Morninggl ry	8	8	7	9	8	8	3	9	7	0	6	8	8	7	6	3	8	6	7	9	8	6	8	6	8	10	7	6	8	8							
Nuts dge	9	9	4	8	10	9	8	3	0	0	9	9	7	3	8	6	9	5	4	8	3	5	10	5	9	9	4	6	10	7							
Rap	9	10	9	8	10	10	10	10	8	6	10	9	10	9	10	5	9	10	7	10	10	10	10	10	10	10	10	9	7	8	10						
Rice Dry Seed	10	9	9	9	10	10	9	9	6	0	4	3	7	4	8	3	8	7	7	9	6	7	10	4	2	4	5	3	3	3							
Sicklepod	8	9	8	4	3	9	3	3	4	0	8	10	9	6	7	3	8	6	6	3	9	4	3	8	4	5	6	7	5	3	0						
S yb an	9	7	9	3	10	8	5	3	3	0	4	3	8	2	5	0	6	6	3	9	4	3	8	4	5	8	6	3	2	5							
Sugar beet	9	9	9	9	9	10	9	9	10	7	9	10	10	7	9	6	8	8	7	10	9	9	9	8	10	8	10	8	8	9							
T aw ed	6	5	6	5	9	9	3	3	-	4	6	7	9	8	9	5	8	5	7	8	5	6	7	5	6	9	8	5	3	3							
V lv tl af	9	9	3	5	9	9	9	9	10	0	9	8	9	7	8	6	9	7	8	10	9	9	9	9	8	10	10	8	10	8	6						
Wh at	8	9	7	4	9	9	5	6	3	0	0	0	6	0	6	0	3	3	2	3	0	0	6	0	0	0	6	0	0	3							
Wild buckwheat	8	9	9	8	9	9	8	8	6	3	6	7	7	6	7	3	8	6	5	8	7	7	8	10	7	8	7	7	6	-							
Wild at	7	7	7	4	9	8	4	3	0	-	0	0	6	0	3	0	7	3	3	3	3	0	0	5	0	0	6	6	0	2	2						
Barnyardgrass	9	10	10	4	10	9	9	9	5	0	6	7	10	6	10	4	9	6	5	10	3	5	10	4	3	10	6	4	2	4							

BNSDOCID: <WO 9115478A1 | >

Table C

Rate (4 g/ha)	COMPOUND																																
	1	2	3	4	5	6	7	8	9	10	11	12	13	15	16	17	18	19	20	22	23	24	25	26	27	28	29	30	31	36			
PREMERGENCE																																	
Barley	5	6	0	0	3	4	0	0	0	0	0	0	4	0	5	0	4	3	0	3	0	0	3	0	0	0	0	4	0	0	2		
Blackgrass	7	5	5	5	10	9	3	3	-	-	-	3	6	0	6	0	7	7	6	3	0	0	0	0	2	2	5	3	0	0	0		
Chickwe d	5	7	6	5	3	8	8	2	7	7	6	7	7	5	8	0	7	7	0	7	7	3	5	0	4	0	6	4	0	3			
Cocklebur	6	8	7	3	9	8	4	2	0	0	7	7	7	3	6	0	10	9	7	7	5	0	7	6	6	6	6	5	7	3			
Corn	3	5	0	0	0	4	2	0	0	0	0	0	6	0	3	0	4	0	3	5	0	0	6	0	5	0	0	0	0	2			
Cotton	6	3	3	0	2	2	0	3	0	0	3	8	7	4	4	0	3	3	0	6	3	4	7	3	3	5	0	2	0	2			
Crabgrass	8	8	7	3	8	9	7	4	8	10	5	7	3	6	8	0	3	3	0	3	4	0	6	0	2	8	3	4	6	3			
Downy br me	5	6	3	0	5	4	3	2	0	0	0	0	3	2	6	0	4	3	0	3	0	0	3	2	0	0	4	0	0	4			
Giant foxtail	7	5	4	3	9	8	5	6	0	0	0	0	3	0	3	0	7	3	0	4	0	0	3	0	2	2	0	0	0	0	0		
Gr en foxtail	7	8	3	3	8	8	2	4	0	0	0	3	3	0	6	0	7	3	0	6	3	0	6	0	0	2	3	0	0	1			
Jimsonwe d	3	3	4	0	8	8	8	2	5	0	6	6	7	7	7	0	4	3	3	8	6	5	8	0	2	0	6	3	4	4			
Johns ngrass	6	3	3	0	3	8	8	3	3	0	3	3	7	6	7	0	8	4	3	7	0	3	7	0	0	5	4	0	3	2			
Lambsquart rs	9	8	9	8	10	9	9	9	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	9	-	-	-	-	7				
Morninggl ry	3	8	7	3	3	3	3	7	6	0	4	7	6	5	0	3	9	4	3	5	5	4	7	5	3	5	6	4	4	4			
Nuts dge	8	7	0	7	9	9	6	0	0	0	3	8	5	0	3	3	9	4	3	8	3	0	7	4	7	2	3	4	4	-			
Rape	9	9	7	7	9	9	10	5	7	0	7	6	10	7	5	0	9	9	7	8	9	6	9	7	3	6	8	3	6	9			
Ric Dry Se d	8	8	8	8	9	9	4	5	4	0	0	0	6	0	7	0	7	5	6	7	3	3	9	2	0	3	3	0	2	2			
Sicklepod	5	3	7	0	3	3	0	0	3	0	7	8	7	5	6	3	0	0	3	8	7	5	10	0	0	5	3	3	0	-			
S ybean	5	2	2	0	8	3	2	0	0	0	0	0	7	0	4	0	4	3	0	4	3	0	6	3	0	2	2	0	0	0	0		
Sugar beet	8	7	8	7	9	9	9	3	7	6	6	7	9	7	8	3	7	7	6	10	7	7	8	8	8	8	8	5	6	6			
Teaweed	3	3	2	0	5	7	0	0	0	3	3	7	8	8	8	3	7	3	6	7	0	5	5	0	2	3	5	4	0	-			
V lvetleaf	9	8	0	3	3	9	7	5	10	-	6	6	7	4	7	4	7	7	8	8	7	7	9	-	4	0	-	9	4	5			
Wh at	5	7	3	0	3	5	2	2	0	0	0	0	3	0	5	0	0	3	0	0	0	0	5	0	0	0	3	0	0	0	0		
Wild buckwheat	8	7	8	7	8	8	8	8	4	0	5	7	7	5	6	0	7	5	4	7	6	7	8	9	4	5	6	3	3	7			
Wild at	6	5	3	2	3	3	2	0	0	-	0	0	3	0	3	0	5	0	0	0	0	0	3	0	0	3	3	0	0	0	0		
Barnyardgrass	6	3	3	0	2	5	5	9	0	0	0	6	8	3	9	0	8	3	3	8	0	0	9	3	3	0	3	0	0	3			



Table C

		COMPOUND																																					
Rate (1 g/ha)		1	2	3	5	6	7	8	9	11	12	13	15	16	17	18	19	20	22	23	24	25	26	27	28	29	30	31	36										
PREEMERGENCE																																							
Barl y	2	3	0	0	2	0	0	0	0	0	0	2	0	3	0	3	0	0	0	0	0	0	0	0	0	0	2	0	0	0	0	0	0	0	0	0			
Blackgrass	5	5	3	3	3	0	0	-	0	0	4	0	4	0	4	0	6	3	3	0	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0		
Chickw ed	5	5	4	0	5	7	0	5	0	6	6	0	7	0	7	0	4	0	0	7	5	0	3	0	0	0	0	3	3	0	0	0	0	0	0	0	0		
Cockl bur	0	3	3	8	2	0	0	0	0	3	6	7	3	5	0	7	7	6	6	3	0	5	0	-	4	-	3	7	0	0	0	0	0	0	0	0	0		
Corn	-	2	0	0	0	0	0	0	0	0	0	2	0	0	0	2	0	2	2	0	0	5	0	4	0	0	0	0	0	0	0	0	0	0	0	0	0		
C tton	3	0	0	0	0	0	0	0	0	0	4	4	4	3	0	2	0	0	5	3	3	6	0	0	3	0	0	0	0	0	0	0	0	0	0	0	0		
Crabgrass	7	5	4	3	5	3	0	8	3	3	0	5	5	0	0	0	0	0	3	0	0	5	0	0	0	0	3	4	3	2	0	0	0	0	0	0	0		
Downy brome	4	3	0	0	2	0	0	0	0	0	0	0	0	3	0	3	0	0	0	0	0	0	0	0	0	0	3	0	0	0	0	0	0	0	0	0	0		
Giant foxtail	4	3	0	5	4	2	4	0	0	0	0	0	0	0	0	4	0	0	0	0	0	3	0	0	0	0	3	0	0	0	0	0	0	0	0	0	0		
Green foxtail	3	4	0	2	4	0	0	0	0	0	2	0	0	0	0	5	0	0	5	3	0	4	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Jims nweed	0	0	0	3	2	3	0	4	0	4	4	4	6	0	5	0	5	0	6	3	3	7	0	0	0	0	0	3	2	3	3	0	0	0	0	0	0		
Johnsongrass	2	3	0	0	2	2	0	0	0	0	6	5	6	0	3	0	0	5	0	6	3	7	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Lambsquarters	5	8	9	7	3	8	5	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	8	-	-	-	-	-	-	-	-	-	-	-	-	-		
M rningglory	0	3	3	0	0	0	3	3	3	3	6	4	6	5	0	3	0	0	4	4	3	6	2	0	3	4	4	3	0	0	0	0	0	0	0	0	0		
Nuts dge	6	5	0	8	3	2	0	0	3	3	4	0	0	0	0	3	3	0	6	3	0	5	3	-	0	3	4	2	3	3	6	5	3	3	3	6	5		
Rap	7	8	7	8	7	7	2	3	3	3	0	5	3	3	0	8	7	5	3	3	0	8	0	3	4	5	0	3	5	3	5	3	3	3	5	3	5		
Ric Dry Seed	8	4	4	3	4	2	2	3	0	0	3	0	6	0	6	3	0	-	5	4	5	8	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Sickl pod	2	0	5	0	0	0	0	0	5	3	5	3	5	0	4	0	2	0	3	-	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Soyb an	0	0	0	2	0	0	0	0	0	0	0	0	0	4	0	2	0	0	3	-	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Sugar beet	7	7	7	9	8	9	0	6	3	5	8	6	4	0	6	6	6	8	5	3	7	0	4	5	5	4	6	2	0	0	0	0	0	0	0	0	0		
Teaweed	0	0	0	0	2	0	0	0	0	0	5	3	8	8	0	3	0	3	3	0	4	3	0	0	-	0	0	0	0	0	0	0	0	0	0	0	0		
Velvetleaf	5	3	0	0	3	3	2	10	3	6	6	3	7	0	5	4	-	7	6	7	8	0	0	-	0	0	0	0	0	0	0	0	0	0	0	0	0		
Wh at	3	3	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Wild buckwh at	7	3	8	2	7	2	3	3	3	5	6	3	5	0	5	4	3	5	5	0	8	3	0	0	0	0	4	3	0	0	0	0	0	0	0	0	0		
Wild oat	4	3	0	0	3	0	0	0	0	0	0	0	0	0	0	4	0	0	0	0	0	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0		
Barnyardgrass	2	0	0	0	2	2	3	0	0	4	7	0	6	0	5	0	0	5	0	0	6	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0		

**TEST D**

5       The compounds evaluated in this test were  
formulated in a non-phytoxic solvent and applied to  
the soil surface before plant seedlings emerged  
(preemergence application), to water that covered the  
soil surface (paddy application), and to plants that  
were in the one-to-four leaf stage (postemergence  
10 application). A sandy loam soil was used for the  
preemergence and postemergence tests, while a silt  
loam soil was used in the paddy test. Water depth  
was approximately 2.5 cm for the paddy test and was  
maintained at this level for the duration of the test.

15       Plant species in the preemergence and  
postemergence tests consisted of barley (Hordeum  
vulgare), bedstraw (Galium aparine), blackgrass  
(Alopecurus myosuroides), chickweed (Stellaria  
media), corn (Zea mays), cotton (Gossypium hirsutum),  
20 crabgrass (Digitaria sanguinalis), downy brome  
(Bromus tectorum), giant foxtail (Setaria faberii),  
lambsquarters (Chenopodium album), morningglory  
(Ipomoea hederacea), pigweed (Amaranthus retroflexus),  
rape (Brassica napus), ryegrass (Lolium multiflorum),  
25 sorghum (Sorghum bicolor), soybean (Glycine max),  
speedwell (Veronica persica), sugar beet (Beta  
vulgaris), velvetleaf (Abutilon theophrasti), wheat  
(Triticum aestivum), wild buckwheat (Polygonum  
convolvulus), and wild oat (Avena fatua). All plant  
30 species were planted one day before application of  
the compound for the preemergence portion of this  
test. Plantings of these species were adjusted to  
produce plants of appropriate size for the  
postemergence portion of the test. Plant species in  
35 the paddy test consisted of barnyardgrass

(Echinochloa crus-galli), rice (Oryza sativa), and umbrella sedge (Cyperus difformis).

5           All plant species were grown using normal  
greenhouse practices. Visual evaluations of injury  
expressed on treated plants, when compared to  
untreated controls, were recorded approximately  
10       fourteen to twenty-one days after application of the  
test compound. Plant response ratings, summarized in  
Table D, were recorded on a 0 to 10 scale where 0 is  
no injury and 10 is complete control. A dash (-)  
response means no test result.

15

20

25

30

35

Tabl D

Rate (62 g/ha)	COMPOUND					
	32	35	40	41	42	43
POSTEMERGENCE						
Barley Igri	9	10	10	5	10	6
Bedstraw	10	10	10	10	10	10
Blackgrass	9	10	10	10	10	10
Chickweed	10	10	10	10	10	10
Corn	10	10	10	9	10	8
Cotton	10	10	10	10	10	10
Crabgrass	7	10	6	0	3	4
Downy brome	9	10	9	6	10	8
Duck salad	-	-	-	9	9	-
Giant foxtail	10	10	10	7	10	9
Lambsquarters	10	10	10	10	10	10
Morningglory	10	10	10	10	10	10
Pigweed	10	10	10	10	10	10
Rape	10	10	10	10	10	10
Ryegrass	10	10	10	3	5	7
Sorghum	10	10	10	7	10	10
Soybean	10	10	10	7	10	9
Speedwell	9	10	10	10	8	9
Sugar beet	10	10	10	10	10	10
Velvetleaf	10	10	10	10	9	10
Wheat	9	9	9	3	9	7
Wild buckwheat	10	10	10	10	9	8
Wild oat	9	9	6	5	4	7
Barnyardgrass	9	10	9	9	10	9
Rice Japonica	8	8	8	8	8	9
Umbrella sedge	10	10	9	9	9	9



Table D

Rate (62 g/ha)	COMPOUND					
	32	35	40	41	42	43
PREEMERGENCE						
Barley Igri	9	5	4	2	9	2
Bedstraw	9	9	10	10	10	10
Blackgrass	9	7	9	7	9	10
Chickweed	10	-	10	9	9	9
Corn	8	10	2	3	7	3
Cotton	6	9	9	10	10	7
Crabgrass	8	10	7	8	5	7
Downy brome	10	7	7	7	10	7
Giant foxtail	7	9	4	6	7	7
Lambsquarters	10	-	10	9	10	10
Morningglory	8	8	10	10	9	8
Pigweed	10	10	10	9	10	9
Rape	10	9	10	10	9	10
Ryegrass	10	10	6	6	7	8
Sorghum	9	10	9	7	10	9
Soybean	7	9	7	0	9	4
Speedwell	10	9	10	9	10	9
Sugar beet	10	9	9	10	10	10
Velvetleaf	8	9	9	8	9	9
Wheat	9	7	4	2	9	4
Wild buckwheat	9	8	9	8	9	9
Wild oat	6	5	5	3	3	3

Tabl D

	COMPOUND											
Rate (31 g/ha)	32	33	34	35	37	38	39	40	41	42	43	
POSTEMERGENCE												
Barley Igri	9	9	9	10	6	10	4	9	4	9	5	
Bedstraw	10	10	10	10	10	10	10	10	10	10	10	
Blackgrass	9	9	9	10	10	10	9	10	9	10	10	
Chickweed	10	10	10	10	10	10	10	10	10	10	10	
Corn	10	10	10	10	8	10	7	10	8	10	7	
Cotton	10	10	10	10	10	10	10	10	10	10	9	
Crabgrass	6	7	10	10	0	8	0	5	0	0	3	
Downy brome	8	10	10	10	5	10	4	9	5	10	7	
Duck salad	-	-	-	-	-	-	-	-	7	8	-	
Giant foxtail	8	10	10	10	6	10	6	10	5	7	8	
Lambsquarters	10	10	10	10	10	10	-	10	10	10	10	
Morningglory	10	8	8	10	10	10	10	10	10	10	10	
Pigweed	10	10	10	10	10	10	0	10	8	10	9	
Rape	10	10	10	10	10	10	10	10	10	10	10	
Ryegrass	9	10	10	10	0	10	4	10	2	4	7	
Sorghum	10	10	10	10	10	10	7	10	6	10	10	
Soybean	10	10	10	10	10	10	10	10	7	10	9	
Speedwell	9	8	8	10	5	8	9	9	10	8	8	
Sugar beet	10	10	10	10	10	10	10	10	10	10	10	
Velvetleaf	10	10	10	10	8	10	10	10	10	9	10	
Wheat	9	9	9	9	4	9	2	9	0	8	7	
Wild buckwheat	10	10	10	10	10	10	9	10	10	9	7	
Wild oat	8	9	10	7	3	9	4	5	4	4	6	
Barnyardgrass	9	8	9	9	8	9	7	9	8	9	9	
Rice Japonica	8	8	9	8	7	9	8	8	8	8	9	
Umbrella sedge	10	9	10	10	9	9	9	9	9	9	9	

Table D

	COMPOUND											
Rate (31 g/ha)	32	33	34	35	37	38	39	40	41	42	43	
PREEMERGENCE												
Barley Igri	8	3	7	4	3	4	0	2	0	7	2	
Bedstraw	8	9	10	9	10	8	9	10	9	10	10	
Blackgrass	9	10	8	6	9	8	7	9	6	9	9	
Chickweed	10	10	10	-	10	9	8	10	9	9	7	
Corn	5	8	9	10	0	6	3	0	0	7	2	
Cotton	6	7	8	9	7	8	3	7	6	9	5	
Crabgrass	8	7	8	10	3	7	0	7	6	4	5	
Downy brome	8	8	10	6	4	8	4	5	6	9	5	
Giant foxtail	6	8	8	9	2	7	10	3	6	6	4	
Lambsquarters	10	10	10	-	10	10	9	10	9	10	9	
Morningglory	8	6	7	7	6	3	6	10	9	9	3	
Pigweed	10	9	9	10	8	8	7	10	9	9	9	
Rape	10	9	9	8	10	10	10	10	10	9	10	
Ryegrass	9	10	10	9	4	5	5	4	6	6	5	
Sorghum	8	9	10	10	5	10	7	9	4	7	9	
Soybean	6	7	2	7	0	6	0	6	0	9	4	
Speedwell	10	10	10	8	9	10	10	10	9	9	9	
Sugar beet	9	9	10	9	10	9	9	9	10	10	9	
Velvetleaf	8	9	7	9	8	8	7	8	8	8	8	
Wheat	6	6	6	6	0	3	0	3	1	5	2	
Wild buckwheat	9	9	8	8	8	9	8	8	7	9	9	
Wild oat	4	4	5	4	0	0	0	3	2	0	0	

Table D

	COMPOUND											
Rate (16 g/ha)	32	33	34	35	37	38	39	40	41	42	43	
POSTEMERGENCE												
Barley Igri	8	9	9	10	5	9	4	8	3	9	4	
Bedstraw	10	10	10	10	10	10	10	10	10	10	9	
Blackgrass	9	9	9	10	10	10	8	9	9	10	10	
Chickweed	10	9	10	10	10	10	10	10	10	10	7	
Corn	10	10	10	10	8	10	6	10	6	10	6	
Cotton	9	10	10	10	10	10	10	10	10	10	9	
Crabgrass	5	6	9	9	0	7	0	4	0	0	0	
Downy brome	7	9	9	8	4	10	4	5	4	10	6	
Duck salad	-	-	-	-	-	-	-	-	1	7	-	
Giant foxtail	7	9	10	10	6	10	5	10	3	6	8	
Lambsquarters	10	10	10	10	10	10	10	10	10	10	9	
Morningglory	10	7	7	10	10	10	10	10	10	10	9	
Pigweed	10	9	10	10	7	9	0	10	7	5	8	
Rape	10	10	10	10	10	10	10	10	10	10	10	
Ryegrass	9	10	10	10	0	10	4	9	0	4	6	
Sorghum	10	10	10	10	10	10	6	10	6	10	10	
Soybean	10	10	10	10	10	10	10	10	7	10	9	
Speedwell	8	8	8	9	5	8	9	8	10	7	7	
Sugar beet	10	10	10	10	10	10	10	10	10	10	8	
Velvetleaf	10	10	10	10	7	10	7	10	10	5	9	
Wheat	6	9	9	4	4	9	0	7	0	8	7	
Wild buckwheat	10	10	9	10	10	10	8	10	10	9	6	
Wild oat	4	8	9	5	2	9	4	4	4	4	5	
Barnyardgrass	8	8	9	9	5	9	5	7	6	9	8	
Rice Japonica	8	8	8	8	7	8	7	8	6	8	8	
Umbrella sedge	9	9	10	9	9	9	5	9	9	9	9	

Table D

	COMPOUND											
Rate (16 g/ha)	32	33	34	35	37	38	39	40	41	42	43	
PREEMERGENCE												
Barley Igri	7	3	6	3	0	2	0	1	0	6	1	
Bedstraw	8	9	10	9	10	8	9	9	9	10	10	
Blackgrass	9	8	7	6	6	7	5	8	5	9	9	
Chickweed	9	10	10	-	10	9	8	9	8	9	4	
Corn	4	8	2	4	0	0	2	0	0	2	0	
Cotton	6	7	8	7	7	7	2	5	-	9	5	
Crabgrass	6	7	7	10	2	5	0	6	6	3	3	
Downy brome	8	7	8	4	0	3	3	4	5	9	2	
Giant foxtail	5	8	7	8	0	6	2	3	6	4	4	
Lambsquarters	10	10	10	-	9	9	8	8	9	9	9	
Morningglory	6	5	3	7	6	3	2	10	7	9	2	
Pigweed	9	9	9	10	8	8	7	10	9	9	8	
Rape	10	9	9	8	9	9	8	9	10	9	10	
Ryegrass	9	10	10	9	0	4	4	4	3	6	5	
Sorghum	3	9	6	10	2	10	4	3	4	4	9	
Soybean	6	6	2	6	0	3	0	5	0	7	2	
Speedwell	10	9	10	8	8	8	9	9	8	9	9	
Sugar beet	9	9	9	9	9	9	9	8	10	10	9	
Velvetleaf	7	8	7	8	8	7	7	7	7	7	7	
Wheat	6	4	4	3	0	0	0	1	0	5	0	
Wild buckwheat	8	8	7	7	8	9	7	7	7	9	8	
Wild oat	0	3	3	3	0	0	0	0	0	0	0	

Table D

	COMPOUND											
Rate (8 g/ha)	32	33	34	35	37	38	39	40	41	42	43	
POSTEMERGENCE												
Barley Igri	8	8	9	9	4	9	3	7	0	9	3	
Bedstraw	10	10	10	10	10	10	10	10	10	10	8	
Blackgrass	9	9	8	10	7	10	6	9	9	10	9	
Chickweed	9	9	10	10	10	10	10	10	10	10	7	
Corn	10	8	10	10	7	9	2	10	2	10	5	
Cotton	9	9	10	10	10	10	10	10	10	10	9	
Crabgrass	0	5	8	7	0	6	0	3	0	0	0	
Downy brome	5	6	9	8	4	9	3	4	3	10	4	
Duck salad	-	-	-	-	-	-	-	-	0	0	-	
Giant foxtail	6	7	10	9	5	10	3	10	0	4	7	
Lambsquarters	10	8	10	9	10	10	10	10	10	10	8	
Morningglory	8	6	7	8	10	10	10	10	10	10	7	
Pigweed	10	8	10	9	7	7	0	10	6	4	8	
Rape	10	10	10	10	10	10	10	10	10	10	10	
Ryegrass	6	9	10	10	0	9	3	7	0	2	5	
Sorghum	7	10	10	10	10	10	6	10	5	9	10	
Soybean	10	10	10	9	10	10	10	10	7	10	9	
Speedwell	8	7	8	9	4	8	9	-	10	6	7	
Sugar beet	10	9	10	10	10	10	10	10	10	10	8	
Velvetleaf	10	10	10	10	7	10	7	10	10	5	9	
Wheat	5	6	9	4	0	8	0	5	0	7	6	
Wild buckwheat	10	8	9	9	9	10	8	10	9	9	6	
Wild oat	4	6	9	3	2	8	3	0	2	2	3	
Barnyardgrass	7	7	9	9	4	7	1	6	4	6	8	
Rice Japonica	8	8	8	8	6	8	1	8	1	6	8	
Umbrella sedge	9	9	9	9	6	9	4	9	7	6	9	

Table D

	COMPOUND											
Rate (8 g/ha)	32	33	34	35	37	38	39	40	41	42	43	
PREEMERGENCE												
Barley Igri	2	0	3	0	0	0	0	0	0	5	0	
Bedstraw	-	9	9	8	7	6	7	7	7	9	9	
Blackgrass	8	8	7	5	4	6	4	7	4	9	6	
Chickweed	9	10	10	-	10	9	7	9	5	9	4	
Corn	0	7	0	2	0	0	0	0	0	0	0	
Cotton	5	7	7	7	4	6	2	4	4	9	4	
Crabgrass	4	4	6	8	0	3	0	6	6	3	2	
Downy brome	6	6	8	3	0	0	0	3	3	5	2	
Giant foxtail	5	3	2	8	0	4	0	3	5	3	2	
Lambsquarters	9	9	9	-	6	7	8	8	7	8	8	
Morningglory	5	4	3	7	3	2	0	9	4	8	0	
Pigweed	9	8	9	10	7	7	7	10	8	9	7	
Rape	9	9	9	8	7	8	5	7	9	9	7	
Ryegrass	4	8	10	7	0	2	2	2	2	4	0	
Sorghum	0	9	4	10	0	4	2	0	3	3	7	
Soybean	4	4	0	5	0	0	0	-	0	6	0	
Speedwell	8	9	10	8	8	7	-	8	8	8	8	
Sugar beet	9	9	9	8	8	8	7	8	9	10	9	
Velvetleaf	6	8	7	8	4	6	3	7	7	7	7	
Wheat	0	3	3	0	0	0	0	0	0	4	0	
Wild buckwheat	7	8	7	7	7	8	6	7	7	8	7	
Wild oat	0	2	2	0	0	0	0	0	0	0	0	

Table D

Rate (4 g/ha)	COMPOUND										
	32	33	34	35	37	38	39	40	41	42	43
POSTEMERGENCE											
Barley Igri	6	7	7	6	4	8	2	5	0	7	0
Bedstraw	9	8	10	10	10	10	10	10	10	10	7
Blackgrass	9	8	8	9	6	9	5	9	0	10	8
Chickweed	9	8	10	10	10	10	10	10	10	10	7
Corn	10	7	7	9	3	8	2	10	0	10	4
Cotton	9	8	10	10	10	10	9	9	10	9	8
Crabgrass	0	4	6	5	0	4	0	2	0	0	0
Downy brome	3	4	8	0	2	4	0	4	0	9	3
Duck salad	-	-	-	-	-	-	-	-	0	0	-
Giant foxtail	5	6	10	8	3	7	3	6	0	2	4
Lambsquarters	10	8	10	8	9	10	10	10	9	10	7
Morningglory	6	4	6	7	10	10	10	10	10	10	3
Pigweed	7	8	10	9	7	7	0	10	5	4	4
Rape	10	10	10	10	9	10	10	10	10	10	10
Ryegrass	5	9	9	10	0	7	0	5	0	0	3
Sorghum	7	9	10	10	7	10	6	10	3	8	10
Soybean	10	10	10	9	10	10	10	10	7	10	9
Speedwell	8	7	7	8	3	8	7	6	9	5	5
Sugar beet	9	8	10	10	10	10	10	10	10	10	8
Velvetleaf	10	10	9	10	7	10	6	10	10	2	9
Wheat	4	4	7	0	0	5	0	3	0	7	4
Wild buckwheat	9	8	9	7	9	10	-	8	9	7	6
Wild oat	2	4	6	0	0	5	2	0	0	0	0
Barnyardgrass	4	5	8	8	0	6	0	2	2	5	5
Rice Japonica	8	8	8	8	2	8	0	7	0	0	7
Umbrella sedge	9	8	9	9	3	9	1	9	3	5	9



Table D

Rate (4 g/ha)	COMPOUND										
	32	33	34	35	37	38	39	40	41	42	43
PREEMERGENCE											
Barley Igri	0	0	0	0	0	0	0	0	0	4	0
Bedstraw	8	8	8	8	6	6	5	7	6	9	9
Blackgrass	7	7	6	4	3	3	3	5	2	8	0.
Chickweed	4	10	10	-	9	8	7	8	5	9	0
Corn	0	6	0	2	0	0	0	0	0	0	0
Cotton	3	7	5	6	2	4	2	3	4	8	3
Crabgrass	4	4	4	7	0	2	0	3	3	3	2
Downy brome	6	4	7	3	0	0	0	3	2	3	0
Giant foxtail	3	3	0	6	0	3	0	0	4	0	0
Lambsquarters	8	9	9	-	6	-	6	8	7	7	8
Morningglory	4	2	2	4	2	0	0	4	0	7	0
Pigweed	9	8	8	10	7	5	4	8	7	7	7
Rape	9	8	8	8	7	8	3	7	7	9	3
Ryegrass	2	6	5	7	0	0	0	0	2	0	0
Sorghum	0	9	4	7	0	2	0	0	3	0	4
Soybean	4	2	0	2	0	0	0	3	0	6	0
Speedwell	8	9	9	8	5	-	9	8	8	8	6
Sugar beet	-	8	8	8	8	8	7	7	7	9	8
Velvetleaf	6	7	6	7	2	5	2	5	6	6	5
Wheat	0	0	0	0	0	0	0	0	0	0	0
Wild buckwheat	7	8	7	7	7	8	5	7	4	8	4
Wild oat	0	0	0	0	0	0	0	0	0	0	0

Table D

Rate (2 g/ha)	COMPOUND				
	33	34	37	38	39
POSTEMERGENCE					
Barley Igri	5	7	3	5	2
Bedstraw	8	10	10	8	8
Blackgrass	7	7	6	8	4
Chickweed	6	9	10	10	8
Corn	7	7	0	7	0
Cotton	5	8	10	8	6
Crabgrass	3	4	0	0	0
Downy brome	2	6	0	3	0
Giant foxtail	5	10	3	5	2
Lambsquarters	7	8	8	10	7
Morningglory	2	4	10	10	10
Pigweed	6	10	0	3	0
Rape	10	10	8	10	10
Ryegrass	6	8	0	4	0
Sorghum	7	9	7	10	6
Soybean	10	9	10	10	10
Speedwell	5	7	3	6	6
Sugar beet	8	10	10	9	10
Velvetleaf	7	7	7	7	6
Wheat	4	6	0	4	0
Wild buckwheat	3	9	7	9	7
Wild oat	3	4	0	4	0
Barnyardgrass	0	-	0	3	0
Rice Japonica	7	7	1	7	0
Umbrella sedge	7	9	3	9	0

09/1911, 072

89

Table D

Rate (2 g/ha)	COMPOUND				
	33	34	37	38	39
PREEMERGENCE					
Barley Igri	0	0	0	0	0
Bedstraw	8	8	5	5	4
Blackgrass	6	5	2	3	3
Chickweed	10	10	9	5	6
Corn	4	0	0	0	0
Cotton	6	5	0	2	0
Crabgrass	2	0	0	0	0
Downy brome	3	6	0	0	0
Giant foxtail	2	0	0	0	0
Lambsquarters	9	9	5	6	5
Morningglory	0	0	0	0	0
Pigweed	7	7	6	5	4
Rape	8	8	6	5	0
Ryegrass	4	7	0	0	0
Sorghum	5	2	0	0	0
Soybean	0	0	0	0	0
Speedwell	9	8	-	4	7
Sugar beet	8	8	6	7	6
Velvetleaf	6	6	0	3	0
Wheat	0	0	0	0	0
Wild buckwheat	7	7	7	6	4
Wild oat	0	0	0	0	0

**TEST E**

Seeds of barnyardgrass (Echinochloa crus-galli), cocklebur (Xanthium pensylvanicum), corn (Zea mays) (soil surface exposed and soil surface covered with perlite), crabgrass (Digitaria spp.), fall panicum (Panicum dichotomiflorum), giant foxtail (Setaria faberii), green foxtail (Setaria viridis), ivyleaf morningglory (Ipomoea hederacea), jimsonweed (Datura stramonium), johnsongrass (Sorghum halepense), ladysthumb smartweed (Polygonum persicaria), lambsquarters (Chenopodium album), redroot pigweed (Amaranthus retroflexus), sorghum (Sorghum bicolor), soybean (Glycine max), and velvetleaf (Abutilon theophrasti) and purple nutsedge (Cyperus rotundus) tubers were planted and treated preemergence with test chemicals dissolved in a non-phytotoxic solvent. These crop and weed species were also treated with postemergence applications of test compounds. Plants ranged in height from two to twenty-five cm for postemergence treatments.

Treated plants and controls were maintained in a greenhouse for approximately 24 days, after which all species were compared to controls and visually evaluated. The ratings, summarized in Table E, are based on a scale of 0 to 10 where 0 is no effect and 10 is complete control. A dash (-) response means no test result.

Table E

	COMPOUND
Rate (500 g/ha)	5
PREEMERGENCE	
Barnyardgrass	10
Cocklebur	10
Corn	6
Crabgrass	10
Fall panicum	10
Giant foxtail	10
Green foxtail	10
Ivlf mrninglory	10
Jimsonweed	10
Johnsongrass	10
Ladysthmb smtwd	10
Lambsquarters	10
Purple nutsedge	10
Redroot pigweed	10
Sorghum	10
Soybean	10
Velvetleaf	10

Table E

Rate (250 g/ha)	COMPOUND				
	5	7	8	9	10
PREEMERGENCE					
Barnyardgrass	10	10	10	10	7
Cocklebur	10	9	6	8	2
Corn	3	9	4	6	0
Crabgrass	9	9	6	8	2
Fall panicum	10	10	7	9	4
Giant foxtail	10	10	10	9	4
Green foxtail	10	10	9	9	3
Ivlf mrninglory	10	9	9	9	3
Jimsonweed	10	4	6	9	2
Johnsongrass	10	10	10	10	6
Ladysthmb smtwd	10	-	-	10	9
Lambsquarters	10	-	-	-	-
Purple nutsedge	10	10	10	10	4
Redroot pigweed	10	10	10	10	10
Sorghum	10	10	10	10	9
Soybean	10	-	-	8	7
Velvetleaf	10	8	8	6	4

Table E

Rate (125 g/ha)	COMPOUND				
	5	7	8	9	10
POSTEMERGENCE					
Barnyardgrass	10	10	10	10	9
Cocklebur	10	10	10	10	10
Corn	7	10	2	6	0
Crabgrass	10	10	7	2	0
Fall panicum	10	10	9	2	0
Giant foxtail	10	10	8	4	2
Green foxtail	10	10	9	4	4
Ivlf mrninglory	10	10	10	10	10
Jimsonweed	10	10	10	10	10
Johnsongrass	10	10	10	10	10
Ladysthmb smtwd	10	-	-	10	6
Lambsquarters	9	10	9	7	6
Perlite corn	5	6	2	4	0
Purple nutsedge	10	10	10	10	4
Redroot pigweed	10	10	10	10	10
Sorghum	10	10	10	10	10
Soybean	10	10	10	10	10
Velvetleaf	10	10	10	10	10

Tabl E

Rate (125 g/ha)	COMPOUND				
	5	7	8	9	10
PREEMERGENCE					
Barnyardgrass	10	10	10	9	4
Cocklebur	8	7	3	6	0
Corn	2	6	2	4	0
Crabgrass	7	7	5	5	0
Fall panicum	10	7	6	6	2
Giant foxtail	10	9	9	7	0
Green foxtail	10	8	8	6	0
Ivlf mrninglory	8	7	7	8	2
Jimsonweed	9	2	3	7	0
Johnsongrass	10	10	10	9	5
Ladysthmb smtwd	10	-	-	10	6
Lambsquarters	10	-	-	-	-
Purple nutsedge	10	10	9	9	0
Redroot pigweed	10	10	10	10	9
Sorghum	10	10	10	9	7
Soybean	9	-	-	6	4
Velvetleaf	9	6	6	3	2



Table E

Rate (64 g/ha)	COMPOUND					
	5	7	8	9	10	24
POSTEMERGENCE						
Barnyardgrass	10	10	10	10	8	9
Cocklebur	10	10	10	10	10	10
Corn	3	9	0	3	0	0
Crabgrass	10	9	5	0	0	0
Fall panicum	10	10	7	0	0	0
Giant foxtail	10	9	7	2	0	0
Green foxtail	10	10	8	2	2	0
Ivlf mrninglory	10	10	9	10	10	10
Jimsonweed	10	7	10	9	10	9
Johnsongrass	10	10	10	10	7	7
Ladysthmb smtwd	10	-	-	10	5	9
Lambsquarters	8	9	8	6	4	7
Perlite corn	2	4	0	2	0	0
Purple nutsedge	10	10	10	10	2	10
Redroot pigweed	8	10	10	10	10	10
Sorghum	10	10	10	10	9	8
Soybean	10	10	10	10	10	10
Velvetleaf	10	10	10	10	10	10

Table E

Rate (64 g/ha)	COMPOUND				
	5	7	8	9	10
PREEMERGENCE					
Barnyardgrass	10	10	8	7	2
Cocklebur	5	7	0	2	0
Corn	0	4	0	3	0
Crabgrass	5	5	3	3	0
Fall panicum	10	6	4	3	0
Giant foxtail	9	6	6	4	0
Green foxtail	10	5	4	3	0
Ivlf mrninglory	6	6	5	5	0
Jimsonweed	6	0	0	4	0
Johnsongrass	10	10	10	7	3
Ladysthmb smtwd	9	-	-	8	4
Lambsquarters	10	-	-	-	-
Purple nutsedge	10	8	8	7	0
Redroot pigweed	10	10	10	10	8
Sorghum	10	10	10	6	5
Soybean	7	-	-	4	2
Velvetleaf	6	5	2	2	0

Table E

Rate (32 g/ha)	COMPOUND					
	5	7	8	9	10	24
POSTEMERGENCE						
Barnyardgrass	10	10	10	10	6	6
Cocklebur	10	10	10	10	10	10
Corn	0	6	0	0	0	0
Crabgrass	6	8	3	0	0	0
Fall panicum	10	8	4	0	0	0
Giant foxtail	10	9	5	0	0	0
Green foxtail	10	10	8	0	0	0
Ivlf mrninglory	10	10	8	9	10	10
Jimsonweed	8	6	8	9	10	8
Johnsongrass	10	10	10	10	6	5
Ladysthmb smtwd	9	-	-	9	5	9
Lambsquarters	7	7	7	4	2	6
Perlite corn	0	3	0	0	0	0
Purple nutsedge	10	10	9	9	2	10
Redroot pigweed	7	10	10	10	10	9
Sorghum	10	10	10	10	8	5
Soybean	10	10	10	10	10	10
Velvetleaf	10	10	9	9	9	10

Table E

Rate (32 g/ha)	COMPOUND				
	5	7	8	9	10
PREEMERGENCE					
Barnyardgrass	8	6	4	3	0
Cocklebur	2	4	0	0	0
Corn	0	2	0	0	0
Crabgrass	3	2	0	0	0
Fall panicum	7	3	2	0	0
Giant foxtail	8	5	2	0	0
Green foxtail	8	4	0	0	0
Ivlf mrninglory	3	3	2	2	0
Jimsonweed	2	0	0	2	0
Johnsongrass	9	7	6	3	0
Ladysthmb smtwd	8	-	-	3	2
Lambsquarters	9	-	-	-	-
Purple nutsedge	9	7	3	3	0
Redroot pigweed	10	9	8	6	6
Sorghum	10	8	5	3	2
Soybean	5	-	-	0	0
Velvetleaf	2	3	0	0	0

Table E

Rate (16 g/ha)	COMPOUND					
	5	7	8	9	10	24
POSTEMERGENCE						
Barnyardgrass	10	10	8	9	4	2
Cocklebur	10	10	9	10	8	10
Corn	0	4	0	0	0	0
Crabgrass	4	6	0	0	0	0
Fall panicum	8	6	3	0	0	0
Giant foxtail	9	8	3	0	0	0
Green foxtail	9	9	7	0	0	0
Ivlf mrninglory	10	9	7	9	9	10
Jimsonweed	7	4	6	7	9	8
Johnsongrass	10	10	9	9	3	2
Ladysthmb smtwd	8	-	-	8	3	7
Lambsquarters	6	6	5	2	0	5
Perlite corn	0	2	0	0	0	0
Purple nutsedge	10	10	6	8	0	7
Redroot pigweed	6	10	9	9	10	7
Sorghum	10	10	10	10	5	3
Soybean	10	10	9	10	10	10
Velvetleaf	10	10	6	7	6	10

100

Table E

Rate (16 g/ha)	COMPOUND	
	5	9
PREEMERGENCE		
Barnyardgrass	6	0
Cocklebur	0	0
Corn	0	0
Crabgrass	0	0
Fall panicum	4	0
Giant foxtail	4	0
Green foxtail	4	0
Ivlf mrninglory	2	0
Jimsonweed	0	0
Johnsongrass	6	0
Ladysthmb smtwd	6	0
Lambsquarters	5	-
Purple nutsedge	7	0
Redroot pigweed	6	3
Sorghum	7	0
Soybean	2	0
Velvetleaf	0	0

101

Table E

Rate (8 g/ha)	COMPOUND					
	5	7	8	9	10	24
POSTEMERGENCE						
Barnyardgrass	10	10	6	7	2	0
Cocklebur	8	8	5	8	7	9
Corn	0	2	0	0	0	0
Crabgrass	2	3	0	0	0	0
Fall panicum	6	3	0	0	0	0
Giant foxtail	8	5	2	0	0	0
Green foxtail	9	8	5	0	0	0
Ivlf mrninglory	9	8	3	8	7	9
Jimsonweed	7	2	3	4	7	7
Johnsongrass	9	10	8	6	2	0
Ladysthmb smtwd	7	-	-	5	2	5
Lambsquarters	3	4	2	0	0	4
Perlite corn	0	0	0	0	0	0
Purple nutsedge	10	9	4	5	0	4
Redroot pigweed	3	9	9	8	9	7
Sorghum	10	10	9	6	3	0
Soybean	10	10	7	10	6	10
Velvetleaf	9	8	3	4	4	8

Table E

Rate (4 g/ha)	COMPOUND					
	5	7	8	9	10	24
POSTEMERGENCE						
Barnyardgrass	9	7	2	3	0	0
Cocklebur	6	5	2	3	3	6
Corn	0	0	0	0	0	0
Crabgrass	0	0	0	0	0	0
Fall panicum	3	0	0	0	0	0
Giant foxtail	5	3	0	0	0	0
Green foxtail	7	6	3	0	0	0
Ivlf mrninglory	7	5	0	3	3	6
Jimsonweed	6	0	0	2	3	4
Johnsongrass	8	9	6	2	0	0
Ladysthmb smtwd	3	-	-	2	0	2
Lambsquarters	0	2	0	0	0	2
Perlite corn	0	0	0	0	0	0
Purple nutsedge	8	6	2	2	0	2
Redroot pigweed	0	8	5	6	6	4
Sorghum	9	9	5	2	0	0
Soybean	10	10	3	8	3	10
Velvetleaf	7	6	0	2	2	5



**TEST F**

Plastic pots were partially filled with silt  
5 loam soil. The soil was then saturated with water.  
Indica and Japonica rice (Oryza sativa) seedlings at  
the 2.0 to 2.5 leaf stage, seeds selected from  
barnyardgrass (Echinochloa crus-galli), bulrush  
(Scirpus mucronatus), duck salad (Heteranthera  
10 Bimosa), umbrella sedge (Cyperus difformis) and  
tubers selected from waterchestnut (Eleocharis spp.),  
were planted into this soil. After planting, water  
levels were raised to 3 cm above the soil surface and  
maintained at this level throughout the test.  
15 Chemical treatments were formulated in a  
non-phytotoxic solvent and applied directly to the  
paddy water. Treated plants and controls were  
maintained in a greenhouse for approximately 21 days,  
after which all species were compared to controls and  
20 visually evaluated. Plant response ratings,  
summarized in Table F, are reported on a 0 to 10  
scale where 0 is no effect and 10 is complete  
control. A dash (-) response means no test result.

25

30

35

104

Table F

	COMPOUND				
Rate (64 g/ha)	27				
PADDY					
Barnyardgrass	6				
Bulrush	9				
Duck salad	10				
Indica rice	2				
Japonica rice	4				
Umbrella sedge	10				
Waterchestnut	10				

	COMPOUND		
Rate (32 g/ha)	11 17 27		
PADDY			
Barnyardgrass	8	5	6
Bulrush	9	9	9
Duck salad	10	10	10
Indica rice	3	0	2
Japonica rice	4	4	4
Umbrella sedge	10	9	10
Waterchestnut	-	-	9

	COMPOUND				
Rate (16 g/ha)	11 17 22 23 27				
PADDY					
Barnyardgrass	7	4	10	7	4
Bulrush	8	9	8	9	7
Duck salad	10	10	10	10	8
Indica rice	3	0	9	4	2
Japonica rice	3	0	9	6	0
Umbrella sedge	10	9	10	10	9
Waterchestnut	-	-	10	8	9

105

Table F

Rate (8 g/ha)	COMPOUND				
	11	17	22	23	27
PADDY					
Barnyardgrass	6	0	9	5	4
Bulrush	7	8	8	6	6
Duck salad	10	4	10	8	7
Indica rice	2	0	8	4	0
Japonica rice	3	0	8	6	0
Umbrella sedge	9	9	10	8	7
Waterchestnut	-	-	6	2	3

Rate (4 g/ha)	COMPOUND				
	11	17	22	23	27
PADDY					
Barnyardgrass	3	0	9	4	3
Bulrush	0	2	8	0	0
Duck salad	9	0	10	3	0
Indica rice	0	0	6	2	0
Japonica rice	0	0	7	4	0
Umbrella sedge	9	7	10	6	5
Waterchestnut	-	-	4	2	3

Rate (2 g/ha)	COMPOUND			
	11	17	22	23
PADDY				
Barnyardgrass	3	0	7	2
Bulrush	0	0	2	0
Duck salad	7	0	9	2
Indica rice	0	0	4	0
Japonica rice	0	0	7	4
Umbrella sedge	7	0	8	3
Waterchestnut	-	-	3	2

106

Table F

Rate (1 g/ha)	COMPOUND	
	22	23
PADDY		
Barnyardgrass	6	0
Bulrush	2	0
Duck salad	3	0
Indica rice	4	0
Japonica rice	6	4
Umbrella sedge	4	2
Waterchestnut	2	2

**TEST G**

Compounds evaluated in this test were

5 formulated in a non-phytoxic solvent and applied to the soil surface before plant seedlings emerged (preemergence application) and to plants that were in the one-to-four leaf stage (postemergence application). A sandy loam soil was used for the

10 preemergence test while a mixture of sandy loam soil and greenhouse potting mix in a 60:40 ratio was used for the postemergence test. Test compounds were applied within approximately one day after planting seeds for the preemergence test. Plantings of these

15 crops and weed species were adjusted to produce plants of appropriate size for the postemergence test. All plant species were grown using normal greenhouse practices. Crop and weed species include winter barley (Hordeum vulgare cv. 'Igri'), bedstraw

20 (Galium aparine), blackgrass (Alopecurus myosuroides), chickweed (Stellaria media), downy brome (Bromus tectorum), field violet (Viola arvensis), green foxtail (Setaria viridis), kochia (Kochia scoparia), lambsquarters (Chenopodium album),

25 Persian speedwell (Veronica persica), rape (Brassica napus cv. 'Jet Neuf'), ryegrass (Lolium multiflorum), sugar beet (Beta vulgaris cv. 'US1'), sunflower (Helianthus annuus cv. 'Russian Giant'), spring wheat (Triticum aestivum cv. 'ERA'), winter wheat (Triticum

30 aestivum cv. 'Talent'), wild buckwheat (Polygonum convolvulus), wild mustard (Sinapis arvensis), wild oat (Avena fatua), and wild radish (Raphanus raphanistrum). Blackgrass and wild oat were treated postemergence at two growth stages. The first stage

35 (1) was when the plants had two to three leaves. The second stage (2) was when the plants had

108

approximately four leaves or in the initial stages of  
tillering. Treated plants and untreated controls  
5 were maintained in a greenhouse for approximately 21  
to 28 days, after which all treated plants were  
compared to untreated controls and visually  
evaluated. Plant response ratings, summarized in  
Table G, are based upon a 0 to 10 scale where 0 is no  
10 effect and 10 is complete control. A dash response  
(-) means no test result.

15

20

25

30

35

109

Table G

	COMPOUND	
Rate (125 g/ha)	37	39
POSTEMERGENCE		
Bedstraw	10	10
Blackgrass (1)	8	6
Blackgrass (2)	8	5
Chickweed	10	10
Downy brome	8	7
Field violet	10	7
Green foxtail	6	4
Persn Speedwell	4	8
Rape	10	10
Ryegrass	2	2
Sugar beet	10	10
Sunflower	10	10
Wheat (Spring)	3	3
Wheat (Winter)	2	3
Wild buckwheat	10	9
Wild mustard	10	10
Wild oat (1)	2	5
Wild oat (2)	0	4
Wild radish	10	10
Winter Barley	3	3

110

Table G

Rate (125 g/ha)	COMPOUND	
	37	39
PREEMERGENCE		
Bedstraw	10	10
Blackgrass (1)	2	0
Blackgrass (2)	2	0
Chickweed	10	9
Downy brome	4	2
Field violet	10	7
Green foxtail	2	2
Persn Speedwell	10	8
Rape	10	10
Ryegrass	4	3
Sugar beet	10	10
Sunflower	7	5
Wheat (Spring)	3	2
Wheat (Winter)	3	2
Wild buckwheat	7	5
Wild mustard	10	10
Wild oat (1)	3	0
Wild oat (2)	2	0
Wild radish	10	9
Winter Barley	4	2



Table G

Rate (64 g/ha)	COMPOUND			
	34	35	37	39
POSTEMERGENCE				
Bedstraw	10	8	10	10
Blackgrass (1)	4	8	6	4
Blackgrass (2)	5	8	5	4
Chickweed	10	10	10	10
Downy brome	9	10	6	5
Field violet	6	5	8	5
Green foxtail	10	9	5	2
Kochia	-	9	-	-
Lambsquarters	10	8	-	-
Persn Speedwell	7	3	3	6
Rape	10	10	10	10
Ryegrass	10	10	0	0
Sugar beet	10	10	10	10
Sunflower	10	10	10	10
Wheat (Spring)	9	10	2	2
Wheat (Winter)	9	10	1	2
Wild buckwheat	10	9	9	8
Wild mustard	10	10	10	10
Wild oat (1)	10	10	0	2
Wild oat (2)	10	10	0	2
Wild radish	10	10	10	10
Winter Barley	10	10	2	2

Table G

Rate (64 g/ha)	COMPOUND			
	34	35	37	39
PREEMERGENCE				
Bedstraw	10	10	10	8
Blackgrass (1)	3	8	0	0
Blackgrass (2)	3	8	0	0
Chickweed	10	10	10	8
Downy brome	6	10	2	0
Field violet	10	10	10	5
Green foxtail	10	10	0	0
Kochia	-	9	-	-
Lambsquarters	-	10	-	-
Persn Speedwell	10	9	10	7
Rape	10	10	10	10
Ryegrass	7	10	2	0
Sugar beet	10	10	10	10
Sunflower	8	10	6	3
Wheat (Spring)	3	9	2	0
Wheat (Winter)	3	8	2	0
Wild buckwheat	4	8	5	3
Wild mustard	10	10	10	10
Wild oat (1)	4	6	0	0
Wild oat (2)	4	6	0	0
Wild radish	9	10	8	7
Winter Barley	5	10	2	0

113

Table G

Rate (32 g/ha)	COMPOUND			
	34	35	37	39
POSTEMERGENCE				
Bedstraw	10	6	8	8
Blackgrass (1)	3	6	4	2
Blackgrass (2)	4	6	4	2
Chickweed	10	10	10	10
Downy brome	8	10	3	3
Field violet	5	3	6	3
Green foxtail	10	9	4	0
Kochia	-	6	-	-
Lambsquarters	10	6	-	-
Persn Speedwell	5	2	2	4
Rape	10	10	10	10
Ryegrass	10	10	0	0
Sugar beet	10	10	10	10
Sunflower	10	10	10	10
Wheat (Spring)	8	8	0	0
Wheat (Winter)	7	7	0	0
Wild buckwheat	8	7	6	6
Wild mustard	10	10	10	10
Wild oat (1)	8	8	0	0
Wild oat (2)	8	8	0	0
Wild radish	10	10	10	10
Winter Barley	10	10	0	0

Table G

Rate (32 g/ha)	COMPOUND			
	34	35	37	39
PREEMERGENCE				
Bedstraw	10	8	7	6
Blackgrass (1)	2	7	0	0
Blackgrass (2)	2	7	0	0
Chickweed	10	10	9	5
Downy brome	5	10	0	0
Field violet	8	10	9	4
Green foxtail	10	10	0	0
Kochia	-	8	-	-
Lambsquarters	-	10	-	-
Persn Speedwell	10	8	8	5
Rape	10	10	10	9
Ryegrass	4	10	0	0
Sugar beet	10	10	10	10
Sunflower	6	9	4	2
Wheat (Spring)	2	7	0	0
Wheat (Winter)	1	6	0	0
Wild buckwheat	3	6	4	2
Wild mustard	10	10	10	9
Wild oat (1)	2	4	0	0
Wild oat (2)	2	5	0	0
Wild radish	7	10	7	4
Winter Barley	4	10	0	0

Table G

Rate (16 g/ha)	COMPOUND			
	34	35	37	39
POSTEMERGENCE				
Bedstraw	8	4	6	5
Blackgrass (1)	2	3	2	0
Blackgrass (2)	2	4	2	0
Chickweed	9	9	10	7
Downy brome	6	8	0	0
Field violet	4	2	3	2
Green foxtail	10	8	2	0
Kochia	-	4	-	-
Lambsquarters	9	3	-	-
Persn Speedwell	3	0	0	2
Rape	10	10	10	10
Ryegrass	10	8	0	0
Sugar beet	9	9	10	10
Sunflower	10	10	10	10
Wheat (Spring)	5	5	0	0
Wheat (Winter)	5	4	0	0
Wild buckwheat	7	3	4	5
Wild mustard	10	10	10	10
Wild oat (1)	7	7	0	0
Wild oat (2)	6	7	0	0
Wild radish	10	10	10	10
Winter Barley	10	9	0	0

116

Table G

Rate (16 g/ha)	COMPOUND			
	34	35	37	39
PREEMERGENCE				
Bedstraw	8	6	5	4
Blackgrass (1)	0	4	0	0
Blackgrass (2)	0	5	0	0
Chickweed	8	8	8	3
Downy brome	4	8	0	0
Field violet	5	10	7	2
Green foxtail	7	8	0	0
Kochia	-	6	-	-
Lambsquarters	-	10	-	-
Persn Speedwell	7	7	6	3
Rape	10	10	8	7
Ryegrass	2	9	0	0
Sugar beet	10	10	9	8
Sunflower	4	7	3	0
Wheat (Spring)	0	4	0	0
Wheat (Winter)	0	4	0	0
Wild buckwheat	2	5	2	0
Wild mustard	10	10	9	8
Wild oat (1)	0	2	0	0
Wild oat (2)	0	2	0	0
Wild radish	6	8	6	3
Winter Barley	2	8	0	0

117

Table G

Rate (8 g/ha)	COMPOUND			
	34	35	37	39
POSTEMERGENCE				
Bedstraw	5	2	4	2
Blackgrass (1)	0	2	0	0
Blackgrass (2)	0	2	0	0
Chickweed	7	8	10	4
Downy brome	4	4	0	0
Field violet	2	0	0	0
Green foxtail	7	5	0	0
Kochia	-	2	-	-
Lambsquarters	7	0	-	-
Persn Speedwell	2	0	0	0
Rape	10	10	10	10
Ryegrass	10	6	0	0
Sugar beet	6	8	10	10
Sunflower	10	10	10	10
Wheat (Spring)	3	3	0	0
Wheat (Winter)	2	2	0	0
Wild buckwheat	4	0	2	2
Wild mustard	10	10	10	8
Wild oat (1)	5	4	0	0
Wild oat (2)	4	5	0	0
Wild radish	10	10	10	10
Winter Barley	8	6	0	0

Table G

Rate (8 g/ha)	COMPOUND			
	34	35	37	39
PREEMERGENCE				
Bedstraw	6	5	3	2
Blackgrass (1)	0	2	0	0
Blackgrass (2)	0	2	0	0
Chickweed	6	7	4	0
Downy brome	2	6	0	0
Field violet	3	9	4	0
Green foxtail	4	6	0	0
Kochia	-	5	-	-
Lambsquarters	-	10	-	-
Persn Speedwell	5	6	3	0
Rape	7	10	6	6
Ryegrass	0	7	0	0
Sugar beet	8	9	7	6
Sunflower	3	6	0	0
Wheat (Spring)	0	3	0	0
Wheat (Winter)	0	2	0	0
Wild buckwheat	0	4	0	0
Wild mustard	8	9	7	6
Wild oat (1)	0	0	0	0
Wild oat (2)	0	0	0	0
Wild radish	4	7	4	0
Winter Barley	0	5	0	0



Table G

Rate (4 g/ha)	COMPOUND			
	34	35	37	39
POSTEMERGENCE				
Bedstraw	3	0	2	0
Blackgrass (1)	0	0	0	0
Blackgrass (2)	0	0	0	0
Chickweed	5	5	10	2
Downy brome	3	2	0	0
Field violet	0	0	0	0
Green foxtail	6	4	0	0
Kochia	-	0	-	-
Lambsquarters	6	0	-	-
Persn Speedwell	0	0	0	0
Rape	10	10	10	10
Ryegrass	7	3	0	0
Sugar beet	4	5	10	8
Sunflower	10	10	10	10
Wheat (Spring)	0	2	0	0
Wheat (Winter)	0	0	0	0
Wild buckwheat	2	0	0	0
Wild mustard	10	10	8	6
Wild oat (1)	3	2	0	0
Wild oat (2)	2	3	0	0
Wild radish	9	10	10	10
Winter Barley	6	3	0	0

120

Table G

Rate (4 g/ha)	COMPOUND			
	34	35	37	39
PREEMERGENCE				
Bedstraw	3	4	2	0
Blackgrass (1)	0	0	0	0
Blackgrass (2)	0	0	0	0
Chickweed	4	4	2	0
Downy brome	0	3	0	0
Field violet	2	6	2	0
Green foxtail	2	4	0	0
Kochia	-	3	-	-
Lambsquarters	-	7	-	-
Persn Speedwell	3	4	0	0
Rape	6	9	4	4
Ryegrass	0	3	0	0
Sugar beet	7	8	6	4
Sunflower	0	5	0	0
Wheat (Spring)	0	0	0	0
Wheat (Winter)	0	0	0	0
Wild buckwheat	0	3	0	0
Wild mustard	4	8	5	3
Wild oat (1)	0	0	0	0
Wild oat (2)	0	0	0	0
Wild radish	2	5	2	0
Winter Barley	0	4	0	0

121

Table G

Rate (2 g/ha)	COMPOUND	
	34	35
POSTEMERGENCE		
Bedstraw	2	0
Blackgrass (1)	0	0
Blackgrass (2)	0	0
Chickweed	4	2
Downy brome	2	0
Field violet	0	0
Green foxtail	3	2
Kochia	-	0
Lambsquarters	5	0
Persn Speedwell	0	0
Rape	10	8
Ryegrass	5	2
Sugar beet	2	3
Sunflower	10	10
Wheat (Spring)	0	0
Wheat (Winter)	0	0
Wild buckwheat	0	0
Wild mustard	10	10
Wild oat (1)	2	0
Wild oat (2)	0	0
Wild radish	8	9
Winter Barley	4	2

122

Table G

Rate (2 g/ha)	COMPOUND	
	34	35
PREEMERGENCE		
Bedstraw	2	3
Blackgrass (1)	0	0
Blackgrass (2)	0	0
Chickweed	2	2
Downy brome	0	0
Field violet	0	4
Green foxtail	0	3
Kochia	-	0
Lambsquarters	-	4
Persn Speedwell	2	2
Rape	4	6
Ryegrass	0	0
Sugar beet	6	6
Sunflower	0	3
Wheat (Spring)	0	0
Wheat (Winter)	0	0
Wild buckwheat	0	2
Wild mustard	2	6
Wild oat (1)	0	0
Wild oat (2)	0	0
Wild radish	0	3
Winter Barley	0	2

123

Table G

	COMPOUND
Rate (1 g/ha)	34
POSTEMERGENCE	
Bedstraw	0
Blackgrass (1)	0
Blackgrass (2)	0
Chickweed	2
Downy brome	0
Field violet	0
Green foxtail	0
Lambsquarters	3
Persn Speedwell	0
Rape	8
Ryegrass	2
Sugar beet	0
Sunflower	10
Wheat (Spring)	0
Wheat (Winter)	0
Wild buckwheat	0
Wild mustard	10
Wild oat (1)	0
Wild oat (2)	0
Wild radish	6
Winter Barley	2

124

Table G

	COMPOUND
Rate (1 g/ha)	34
PREEMERGENCE	
Bedstraw	0
Blackgrass (1)	0
Blackgrass (2)	0
Chickweed	0
Downy brome	0
Field violet	0
Green foxtail	0
Lambsquarters	-
Persn Speedwell	0
Rape	3
Ryegrass	0
Sugar beet	4
Sunflower	0
Wheat (Spring)	0
Wheat (Winter)	0
Wild buckwheat	0
Wild mustard	0
Wild oat (1)	0
Wild oat (2)	0
Wild radish	0
Winter Barley	0

**TEST H**

Seeds of alfalfa (Medicago sativa), bean  
5 (Phaseolus vulgaris), bluegrass (Poa pratensis),  
cabbage (Brassica rapa), carrot (Daucus sativa), corn  
(Zea mays), flax (Linum usitatissimum), lettuce  
(Lactuca sativa), lupine (Lupinus albus), oats (Avena  
10 sativa), onion (Allium cepa), pea (sativum), peanut  
(Arachis hypogaea), potato (Solanum tuberosum), rye  
(Secale cereal), sorghum (Sorghum bicolor), sunflower  
(Helianthus annuus), and tomato (Lycopersicon  
esculentum) were planted and treated preemergence  
15 with a test chemical dissolved in a non-phytotoxic  
solvent. These crop species were also treated with  
postemergence applications of the test chemical.  
Plants ranged in height from four to twenty cm (two  
to three leaf stage) when post-emergence applications  
20 were applied. Treated plants and controls were grown  
under greenhouse conditions for approximately  
twenty-four days, after which all plants treated with  
the test chemical were compared to untreated controls  
and visually evaluated for injury response.  
Application rates for the test chemical are shown in  
25 Table H. Plant response ratings, summarized in Table  
H, are from 0 to 10 where 0 is no injury and 10 is  
complete control. A dash (-) response means no test  
result.

30

35

126

Table H

	COMPOUND
Rate (250 g/ha)	5
PREEMERGENCE	
Alfalfa	9
Bean	9
Bluegrass	10
Cabbage	10
Carrott	9
Corn	8
Flax	10
Lettuce	9
Lupine	10
Oats	9
Onion	10
Pea	10
Peanut	9
Potato	10
Rye	10
Sorghum	10
Sunflower	10
Tomato	9



127

Table H

	COMPOUND
Rate (125 g/ha)	5
POSTEMERGENCE	
Alfalfa	9
Bean	10
Bluegrass	6
Cabbage	9
Carrott	10
Corn	8
Flax	9
Lettuce	9
Lupine	10
Oats	9
Onion	8
Pea	7
Peanut	9
Potato	7
Rye	9
Sorghum	10
Sunflower	10
Tomato	9

128

Table H

	COMPOUND
Rate (125 g/ha)	5
PREEMERGENCE	
Alfalfa	9
Bean	9
Bluegrass	9
Cabbage	10
Carrott	9
Corn	8
Flax	10
Lettuce	9
Lupine	10
Oats	9
Onion	10
Pea	10
Peanut	9
Potato	9
Rye	10
Sorghum	10
Sunflower	10
Tomato	9

129

Table H

	COMPOUND
Rate (64 g/ha)	5
POSTEMERGENCE	
Alfalfa	8
Bean	10
Bluegrass	6
Cabbage	9
Carrott	9
Corn	7
Flax	6
Lettuce	9
Lupine	10
Oats	7
Onion	7
Pea	6
Peanut	8
Potato	7
Rye	9
Sorghum	9
Sunflower	10
Tomato	8

130

Table H

	COMPOUND
Rate (64 g/ha)	5
PREEMERGENCE	
Alfalfa	9
Bean	9
Bluegrass	9
Cabbage	10
Carrott	8
Corn	5
Flax	9
Lettuce	10
Lupine	8
Oats	9
Onion	10
Pea	10
Peanut	9
Potato	9
Rye	10
Sorghum	10
Sunflower	9
Tomato	9

131

Table H

## COMPOUND

Rate (32 g/ha)	5
POSTEMERGENCE	
Alfalfa	8
Bean	9
Bluegrass	7
Cabbage	6
Carrott	9
Corn	3
Flax	6
Lettuce	10
Lupine	10
Oats	7
Onion	7
Pea	5
Peanut	9
Potato	7
Rye	9
Sorghum	9
Sunflower	10
Tomato	7

132

Table H

	COMPOUND
Rate (32 g/ha)	5
PREEMERGENCE	
Alfalfa	7
Bean	9
Bluegrass	8
Cabbage	9
Carrott	7
Corn	3
Flax	7
Lettuce	7
Lupine	6
Oats	9
Onion	9
Pea	10
Peanut	7
Potato	8
Rye	9
Sorghum	10
Sunflower	9
Tomato	9

133

Table H

	COMPOUND
Rate (16 g/ha)	5
POSTEMERGENCE	
Alfalfa	6
Bean	9
Bluegrass	6
Cabbage	7
Carrott	9
Corn	1
Flax	6
Lettuce	9
Lupine	10
Oats	8
Onion	6
Pea	6
Peanut	7
Potato	4
Rye	7
Sorghum	9
Sunflower	10
Tomato	4

134

Table H

	COMPOUND
Rate (16 g/ha)	5
PREEMERGENCE	
Alfalfa	6
Bean	2
Bluegrass	6
Cabbage	9
Carrott	2
Corn	1
Flax	6
Lettuce	3
Lupine	8
Oats	9
Onion	7
Pea	10
Peanut	6
Potato	6
Rye	9
Sorghum	9
Sunflower	9
Tomato	8



135

Table H

	COMPOUND
Rate (8 g/ha)	5
POSTEMERGENCE	
Alfalfa	3
Bean	8
Bluegrass	6
Cabbage	2
Carrott	8
Corn	0
Flax	3
Lettuce	9
Lupine	10
Oats	7
Onion	6
Pea	6
Peanut	6
Potato	2
Rye	7
Sorghum	9
Sunflower	10
Tomato	3

136

CLAIMS


5

What is claimed is:

1. A compound selected from

10

15



I

20

wherein:

A is OR<sup>1</sup> or N(CH<sub>3</sub>)<sub>2</sub>;R<sup>1</sup> is C<sub>1</sub>-C<sub>3</sub> alkyl, CH<sub>2</sub>CH=CH<sub>2</sub>, CH<sub>2</sub>C≡CH, CH<sub>2</sub>CH<sub>2</sub>Cl  
or CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>;

25

R<sup>2</sup> is CH<sub>2</sub>F, CHF<sub>2</sub>, CHFCH<sub>3</sub> or CH<sub>2</sub>CN;X is CH<sub>3</sub> or OCH<sub>3</sub>; andY is H, Cl, CH<sub>3</sub>, C<sub>2</sub>H<sub>5</sub>, OCH<sub>3</sub> or OCF<sub>2</sub>H;and their agriculturally suitable salts; provided  
that when Y is Cl, then X is OCH<sub>3</sub>.

30

2. A compound of Claim 1 wherein

A is OR<sup>1</sup>; andR<sup>1</sup> is CH<sub>3</sub>, CH<sub>2</sub>CH<sub>3</sub> or CH(CH<sub>3</sub>)<sub>2</sub>.

35

3. A compound of Claim 2 where


when one of X and Y is CH<sub>3</sub>, then theother of X and Y is other than OCH<sub>3</sub>.

4. The compound of Claim 1,  
methyl 3-(cyanomethyl)-2-[[[(4,6-  
5 dimethyl-2-pyrimidinyl)amino]carbonyl]-  
amino]sulfonyl]benzoate.
5. The compound of Claim 1,  
methyl 2-[[[(4-chloro-6-methoxy-2-  
10 pyrimidinyl)amino]carbonyl]amino]-  
sulfonyl]-3-(cyanomethyl)benzoate.
6. The compound of Claim 1,  
methyl 3-(cyanomethyl)-2-[[[(4-  
15 (difluoromethoxy)-6-methoxy-2-pyrimi-  
dinyl)amino]carbonyl]amino]sulfonyl]-  
benzoate.
7. The compound of Claim 1,  
20 methyl 3-(difluoromethyl)-2-[[[(4-  
methoxy-2-pyrimidinyl)amino]carbonyl]-  
amino]sulfonyl]benzoate.
8. The compound of Claim 1,  
25 methyl 3-(fluoromethyl)-2-[[[(4-  
methoxy-2-pyrimidinyl)amino]carbonyl]-  
amino]sulfonyl]benzoate.
9. The compound of Claim 1,  
30 methyl 2-[[[(4,6-dimethoxy-2-pyrimi-  
dinyl)amino]carbonyl]amino]sulfonyl]-3-  
(fluoromethyl)benzoate.

138

10. The compound of Claim 1,  
ethyl 2-[[[(4-chloro-6-methoxy-2-  
pyrimidinyl)amino]carbonyl]amino]-  
sulfonyl]-3-(fluoromethyl)benzoate.

11. A compound selected from



II

wherein:

A is  $\text{OR}^1$  or  $\text{N}(\text{CH}_3)_2$ ;

$\text{R}^1$  is  $\text{C}_1$ - $\text{C}_3$  alkyl,  $\text{CH}_2\text{CH}=\text{CH}_2$ ,  $\text{CH}_2\text{C}\equiv\text{CH}$ ,  
 $\text{CH}_2\text{CH}_2\text{Cl}$  or  $\text{CH}_2\text{CH}_2\text{OCH}_3$ ;

$\text{R}^2$  is  $\text{CH}_2\text{F}$ ,  $\text{CHF}_2$ ,  $\text{CHFCH}_3$  or  $\text{CH}_2\text{CN}$ ; and

$\text{Z}^1$  is  $\text{Cl}$  or  $\text{NHSiR}^3\text{R}^4\text{R}^5$ ;

$\text{R}^3$  is  $\text{C}_1$ - $\text{C}_4$  alkyl;

$\text{R}^4$  is  $\text{C}_1$ - $\text{C}_4$  alkyl; and

$\text{R}^5$  is  $\text{C}_1$ - $\text{C}_4$  alkyl.

12. A compound of Claim 11 wherein A is  $\text{OR}^1$   
and  $\text{R}^1$  is  $\text{C}_1$ - $\text{C}_2$  alkyl.

13. A compound of Claim 12 wherein  $\text{R}^3$  and  $\text{R}^4$   
are  $\text{CH}_3$  and  $\text{R}^5$  is  $\text{C}(\text{CH}_3)_3$ .

14. A composition suitable for controlling the growth of undesired vegetation which comprises an effective amount of a compound of Claim 1 and at least one of the following: surfactant, solid diluent or liquid diluent.

15. A composition suitable for controlling the growth of undesired vegetation which comprises an effective amount of a compound of Claim 2 and at least one of the following: surfactant, solid diluent or liquid diluent.

16. A composition suitable for controlling the growth of undesired vegetation which comprises an effective amount of a compound of Claim 3 and at least one of the following: surfactant, solid diluent or liquid diluent.

17. A composition suitable for controlling the growth of undesired vegetation which comprises an effective amount of a compound of Claim 4 and at least one of the following: surfactant, solid diluent or liquid diluent.

18. A composition suitable for controlling the growth of undesired vegetation which comprises an effective amount of a compound of Claim 5 and at least one of the following: surfactant, solid diluent or liquid diluent.

19. A method for controlling the growth of undesired vegetation which comprises applying to the locus to be protected an effective amount of the compound of Claim 1.

20. A method for controlling the growth of  
5 undesired vegetation which comprises applying to the  
locus to be protected an effective amount of the com-  
pound of Claim 2.

21. A method for controlling the growth of  
10 undesired vegetation which comprises applying to the  
locus to be protected an effective amount of the com-  
pound of Claim 3.

22. A method for controlling the growth of  
15 undesired vegetation which comprises applying to the  
locus to be protected an effective amount of the com-  
pound of Claim 4.

23. A method for controlling the growth of  
20 undesired vegetation which comprises applying to the  
locus to be protected an effective amount of the com-  
pound of Claim 5.

24. The method of Claim 22 wherein the locus  
25 to be protected is corn.

25. A method for controlling the growth of  
undesired vegetation in wheat or barley which  
comprises applying to the wheat or barley an  
30 effective amount of the compound of Claim 9.

## INTERNATIONAL SEARCH REP RT

International Application No

PCT/US 91/01075

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (If several classification symbols apply, indicate all) *		
According to International Patent Classification (IPC) or to both National Classification and IPC		
IPC <sup>5</sup> : C 07 D 239/42, C 07 D 239/47, C 07 D 239/52, C 07 C 309/89, A 01 N 43/54		
<b>II. FIELDS SEARCHED</b>		
Minimum Documentation Searched *		
Classification System	Classification Symbols	
IPC <sup>5</sup>	C 07 D 239/00, C 07 D 251/00, C 07 C 309/00, A 01 N 43/00	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched *		
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT</b> *		
Category *	Citation of Document, ** with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>
A	EP, A2, 0 096 002 (CIBA-GEIGY) 07 December 1983 (07.12.83), see claims 1, 10, 11, 12, 17.	1, 14, 19, 24, 25
A	EP, A2, 0 073 627 (DU PONT) 09 March 1983 (09.03.83) see claims 1, 25, 26.	1, 14, 19
X	WO, A1, 89/09 214 (DU PONT) 05 October 1989 (05.10.89), see examples. see claims 1, 8, 15.	11
A		1, 14, 19
A	EP, A1, 0 181 526 (CHUGAI) 21 May 1986 (21.05.86), see claim 1.	11
<p>* Special categories of cited documents: <sup>14</sup></p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"Z" document member of the same patent family</p>		
<b>IV. CERTIFICATION</b>		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
15 April 1991	10.06.91	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	Nuria TORIBIO	

Form PCT/ISA/210 (second sheet) (January 1985)

ANHANG  
zum internationalen Recherchen-  
bericht über die internationale  
Patentansmeldung Nr.

ANNEX  
to the International Search  
Report to the International Patent  
Application No.

ANNEXE  
au rapport de recherche inter-  
national relatif à la demande de brevet  
international n°

PCT/US91/01075 SAE 45036

In diesem Anhang sind die Mitglieder  
der Patentfamilien der im obenge-  
nannten internationalen Recherchenbericht  
angeführten Patentdokumente angegeben.  
Diese Angaben dienen nur zur Unter-  
richtung und erfolgen ohne Gewähr.

This Annex lists the patent family  
members relating to the patent documents  
cited in the above-mentioned inter-  
national search report. The Office is  
in no way liable for these particulars  
which are given merely for the purpose  
of information.

La présente annexe indique les  
membres de la famille de brevets  
relatifs aux documents de brevets cités  
dans le rapport de recherche inter-  
national visée ci-dessus. Les renseigne-  
ments fournis sont donnés à titre indica-  
tif et n'engagent pas la responsabilité  
de l'Office.

Im Recherchenbericht angeführtes Patentdokument Patent document cited in search report Document de brevet cité dans le rapport de recherche	Datum der Veröffentlichung Publication date Date de publication	Mitglied(er) der Patentfamilie Patent family member(s) Membre(s) de la famille de brevets	Datum der Veröffentlichung Publication date Date de publication
EP-A1- 181526	21-05-86	AT-E - 40112	15-02-89
		CA-A1- 1255329	06-06-89
		DE-CO- 3567643	23-02-89
		EP-B1- 181526	18-01-89
		JP-A2-61097257	15-05-86
		US-A - 4698364	06-10-87
		JP-A2-61097219	15-05-86
EP-A2- 73627	09-03-83	US-A - 4491467	01-01-85
		AT-E - 23999	15-12-86
		AU-A1-87517/82	03-03-83
		AU-B2- 555081	11-09-86
		BR-A - 8204883	02-08-83
		CA-A1- 1220203	07-04-87
		DE-CO- 3274547	15-01-87
		DK-A - 3754/82	25-02-83
		EP-A3- 73627	28-03-84
		EP-B1- 73627	03-12-86
		ES-A1- 515183	16-10-83
		ES-A1- 8406072	16-10-84
		IL-AO- 66617	31-12-82
		IL-A1- 66617	31-03-86
		JP-A2-58043959	14-03-83
		NZ-A - 201676	31-05-85
		PL-A1- 238013	09-05-83
		US-A - 4555261	26-11-85
		ZA-A - 8206115	28-03-84
EP-A2- 96002	07-12-83	AT-E - 28637	15-08-87
		CA-A1- 1228355	20-10-87
		DE-CO- 3372760	03-09-87
		EP-A3- 96002	25-07-84
		EP-B1- 96002	29-07-87
		IL-AO- 69770	30-09-83
		IL-A1- 68770	29-04-86
		JP-A2-58216167	15-12-83
		US-A - 4537618	27-08-85
		US-A - 4594097	10-06-86
		US-A - 4727186	23-02-88
		ZA-A - 8303779	29-02-84
		IL-AO- 69770	30-12-83
		AU-A1-19566/83	05-04-84
		DD-A5- 207714	14-03-84
		DK-AO- 4389/83	26-09-83
		EP-A2- 111993	27-06-84
		EP-A3- 111993	25-07-84
		FI-AO- 833424	23-09-83
		FI-A - 833424	28-03-84
		GB-AO- 8325360	26-10-83
		GB-A1- 2127408	11-04-84
		GB-B2- 2127408	14-05-86
		IL-AO- 69770	30-12-83
		JP-A2-59080666	10-05-84
		PL-A1- 243905	30-01-85
		PT-A - 77398	01-10-83
		PT-B - 77398	20-03-86
		US-A - 4492708	08-01-85
		ZA-A - 8306958	24-04-85



WD-A1- 8909214

05-10-89

AU-A1-34300/89

16-10-89

DK-A0- 2288/90

21-09-90

DK-A - 2288/90

23-11-90

EP-A1- 336587

11-10-89

EP-A1- 406322

09-01-91

**THIS PAGE BLANK (USPTO)**